



Modulation of Short-Chain Fatty Acid Levels in Chick Gut Using Lactobacillus-Fermented Chicken Feather and Bean Chaff: Implications for Poultry Health

Ezeoke, F. C.¹, Iheukwumere, I. H.², Iheukwumere, C. M.³, Unaeze, B. C.⁴, Ezendianefo, J. N.⁵, Abba, O.⁶, Idigo, M. A.⁷, Aniekwu, C. C.³, Anagor, I. S.⁸, Ochibulu, S. C.² and Nnadozie, H. C.²

¹Department of Animal Science, Faculty of Agriculture, Chukwuemeka Odumegwu Ojukwu University, Uli, Campus, Anambra State, Nigeria

²Department of Microbiology, Faculty of Natural Sciences, Chukwuemeka Odumegwu Ojukwu University, Uli, Campus, Anambra State, Nigeria

³Department of Applied Microbiology and Brewing, Faculty of Biosciences, Nnamdi Azikiwe University, Awka, Nigeria

⁴Department of Microbiology, University of Agriculture and Environmental Sciences, Umuagwo, Imo State, Nigeria



⁵Department of Microbiology, Tansian University, Umunya, Anambra State, Nigeria

⁶Department of Microbiology, Federal University of Gusau, Zamfara State

⁷Department of Biological Sciences, Faculty of Natural Sciences, Chukwuemeka Odumegwu Ojukwu University, Uli, Campus, Anambra State, Nigeria.

⁸Department of Medical Microbiology and Parasitology, Faculty of Basic and Clinical Sciences, Nnamdi Azikiwe University, Nnewi Campus.

*Corresponding author email: fc.ezeoke@coou.edu.ng / ik.iheukwumere@coou.edu.ng

Abstract	Article History
<p>Maintaining healthy gut function in chicks remains a challenge for the poultry industry, impacting growth and productivity. Encouragingly, Lactobacillus-fermented feed additives have shown efficacy in promoting gut health and alleviating these issues. This study aimed to investigate the effects of Lactobacillus acidophilus-fermented chicken feather and bean chaff on modulating short-chain fatty acids (SCFAs) in chick gut. The fermenter was isolated and characterized using cultural, morphological, and biochemical tests, and authenticated using molecular techniques. The isolate was identified as Lactobacillus acidophilus strain DSM20079. The SCFAs profile of the control and test groups were analyzed using gas chromatography. The results showed that the test group had increased levels of acetate ($70.35\% \pm 0.12$ vs $66.22\% \pm 0.11$), propionate ($29.40\% \pm 0.22$ vs $22.40\% \pm 0.16$), and n-valerate ($3.17\% \pm 0.01$ vs $3.03\% \pm 0.01$) compared to the control group, while n-butyrate decreased ($0.97 \text{ mg/L} \pm 0.01$ vs $2.06 \text{ mg/L} \pm 0.01$). Statistical analysis revealed significant differences ($p < 0.05$) in the levels of acetate, n-butyrate, propionate, and n-valerate between the test and control groups. The study concludes that Lactobacillus acidophilus-fermented chicken feather and bean chaff modulates SCFAs and has the potential to enhance gut health in chicks. This study contributes to knowledge by providing evidence on the prebiotic potential of Lactobacillus acidophilus-fermented chicken feather and bean chaff in modulating SCFAs and promoting gut health in poultry.</p> <p>Keywords: <i>Lactobacillus acidophilus</i>, Chicken feather, Bean chaff, Short-chain fatty acids, Gut health, Prebiotic</p>	<p>Received: 29 Dec 2025 Accepted: 08 Feb 2026 Published: 13 Feb 2026</p> <p>Scan QR code to view*</p>  <p>License: CC BY 4.0</p>  <p>Open Access article.</p>
<p>How to cite this paper: Ezeoke, F. C., Iheukwumere, I. H., Iheukwumere, C. M., Unaeze, B. C., Ezendianefo, J. N., Abba, O., Idigo, M. A., Aniekwu, C. C., Anagor, I. S., Ochibulu, S. C., & Nnadozie, H. C. (2026). Modulation of Short-Chain Fatty Acid Levels in Chick Gut Using Lactobacillus-Fermented Chicken Feather and Bean Chaff: Implications for Poultry Health. <i>IPS Interdisciplinary Journal of Biological Sciences</i>, 6(1), 247–256. https://doi.org/10.54117/ijbs.v6i1.123</p>	

Introduction

Normal microbiota of the gut play a crucial role in recycling nutrients that living organisms require for optimum physiology and anatomy (Awad *et al.*, 2017; Okeke *et al.*, 2017; Dim *et al.*, 2025a). Short-chain fatty acid is an essential component found in the gut of living organisms (Awad *et al.*, 2017). This compound contains mainly fat and acid and confers enormous benefits to the organism. The fat and acid components appear in chains, which do not elongate. The roles of short-chain fatty acids involve regulating the integrity of the gut, stabilizing immune function for wading off pathogens, and minimizing inflammation (Latorre *et al.*, 2018). Poor

production of short-chain fatty acid in the gut leads to destabilization of body function in both animals and humans, due to loss of coordination (Gharib-Naseri *et al.*, 2019; Amadi *et al.*, 2017; Dim *et al.*, 2025b).

It is worthy to not that short chain fatty acid is produced naturally in the gut by normal flora, which are vital bacterial species that are capable of digesting dietary fibre (Papatsiros *et al.*, 2013). Digestion of dietary fibre by the normal flora of the gut releases short chain fatty acids as the byproduct such as butyrate, propionate, and acetate (Correa-Oliveira *et al.*,

2016). This indicates that enhancing short chain fatty acid production in both animal and man is essential.

Research has revealed that *Lactobacillus* species are capable of enhancing the production of short chain fatty acid in animals such as chick (Dittoe *et al.*, 2018). *Lactobacillus* species are lactic acid bacteria that are capable of fermenting dietary fibre, which resists the action of digestive enzymes (Papatsiros *et al.*, 2013; Dim *et al.*, 2025c; Chude *et al.*, 2020). Naturally, indigestible fibre can be found in both plant and animals such as bark of maize and beans chaff. The action of the lactic acid and other metabolites that are produced by *Lactobacillus* species facilitates the breakdown of the indigestible substances to release short chain fatty acids (Khan and Iqbal, 2016). Regulation of the intestinal mucosa of birds is highly essential to optimum growth and productivity (Timbermont *et al.*, 2010) and the product enhances absorption of nutrients and stimulation of vibrant immune response in chicks (Stefanello *et al.*, 2020).

Several researchers have worked on the role of organic acid in modulating the integrity of short chain fatty acid such as Awad *et al.* (2017), Latorre *et al.* (2018), and Gharib-Naseri *et al.* (2019) but few studies are available on modulation of short chain fatty acid level of chicks gut using *Lactobacillus* fermented chicken feather and beans chaff. Hence, this study aims to evaluate modulation of short chain fatty acid level of chicks gut using *Lactobacillus* fermented chicken feather and beans chaff.

Materials and Methods

Isolation of the Test Sample

The media used for this isolation was de Man Rogosa and Sharpe broth (MRS) (BIOTECH). A 1.0 ml of fermented yoghurt (Aqua yoghurt) and banana extract were aseptically introduced into sterile Petri dishes (90 mm x 15 mm), then 20 ml of MRS which was prepared according to the manufacturers instruction and the procedures described in Cheesbrough (2010) was added into the plates, allowed to solidified. The plates were incubated in a microaerophilic environment (containing candle used to evacuate all traces of oxygen thereby creating an environment having only carbon iv oxide). The incubation was done for 24 – 48 h at (30±2⁰C). This was carried out using the method described by Egbe *et al.* (2025b), Egbe *et al.* (2025c), Iheukwumere *et al.* (2025d), Iheukwumere *et al.* (2025e).

Characterization and identification of the isolates

The isolates were subcultured on nutrient agar (Biotech), incubated in an inverted position at 37±2⁰C for 24 h. The isolates were characterized and identified using their colonial and morphological descriptions as described in the study published by Iheukwumere *et al.* (2018b), Iheukwumere *et al.* (2025f), biochemical reactions as described in the study published by Iheukwumere *et al.* (2020a), Iheukwumere *et al.* (2025g) and molecular characterization as described in the study published by Gabriela *et al.* (2014), Ekesiobi *et al.* (2025), Ekechukwu *et al.* (2025a), Ekechukwu *et al.* (2025b), Ezedianafo *et al.* (2025a), and Ezedianafo *et al.* (2025b).

Morphological characteristics of the isolates: The cultural descriptions (size, appearance, edge, elevation, and colour) of

the isolates were carried out. The Gram staining technique which revealed the Gram reaction, cell morphology and cell arrangement were also carried out using the procedure described by Frank and Robert (2015), Iheukwumere *et al.* (2020b), Idigo *et al.* (2025a), Idigo *et al.* (2025b), Idigo *et al.* (2025c), Idigo *et al.* (2025d), and Ezedianafo *et al.* (2025c).

Gram staining technique: A thin smear was made on a cleaned, grease-free microscopic slide (75 mm × 25 mm), air-dried, and heat-fixed (Ejike *et al.*, 2017; Iheukwumere *et al.*, 2017a; Iheukwumere *et al.*, 2017b; Iheukwumere *et al.*, 2023a; Iheukwumere *et al.*, 2023b). The smear was flooded with crystal violet solution (0.2%) for 60 seconds and rinsed with clean water. Gram iodine solution (0.01%) was then applied and allowed for 60 seconds. This was rinsed with clean water. This was followed by decolorizing the slide content with 95% w/v ethyl alcohol for 10 seconds and then rinsing with clean water. The smear was then counterstained with safranin solution (0.025%) for 60 seconds, rinsed with cleaned water, blot drained, and air dried. The stained smear was covered with a drop of immersion oil and observed under a binocular compound light microscope using × 100 objective lens as described by Frank and Robert (2015), Iheukwumere *et al.* (2017c), Iheukwumere *et al.* (2018c) Ike *et al.* (2025a), Iheukwumere *et al.* (2024).

Motility test: A semi-solid medium prepared by mixing 5.0 g of bacteriological agar (BIOTECH) with 2.0 g of nutrient broth (BIOTECH) in 1 Litre of distilled water was used. The solution was dissolved and sterilized using autoclaving technique after dispensing 10ml portion in different test tubes. The test tubes were allowed to set in vertical positions and then inoculate the test organisms by performing a single stab down the centre of the test tube to half the depth of the medium using sterile stabbing needle. The test tubes were kept in an incubator in vertical position at 35±2⁰C for 24 h as described by Frank and Robert (2015), Iheukwumere *et al.* (2017d), Iheukwumere *et al.* (2022b), Iheukwumere *et al.* (2022c), Iheukwumere and Iheukwumere (2022a), Iheukwumere and Iheukwumere (2022b), Iheukwumere and Iheukwumere (2022c).

Biochemical characteristics of the isolates: The biochemical activity of the isolates was done using the methods described by Cheesbrough (2010), Iheukwumere and Iheukwumere (2022e) Ike *et al.* (2025b) Ike *et al.* (2025c) Iheukwumere *et al.* (2022d), Idigo *et al.* (2025e), Obiefuna *et al.* (2025a).

Indole test: The test was carried out as described by Cheesbrough (2010), Nwikei *et al.* (2017), Obianom *et al.* (2024), Ekechukwu *et al.* (2025c), Obiefuna *et al.* (2025b), Iheukwumere and Iheukwumere (2022g), and Iheukwumere *et al.* (2022f). Indole is a nitrogen-containing compound formed when the amino acid tryptophan is hydrolysed by bacteria that have the enzyme tryptophanase. This is detected by using KOVAC's reagent. For this test, isolates were cultured in peptone water in 500.0 mL of deionized water. Ten millilitres of peptone water was dispensed into the test tubes and sterilized. The medium was then inoculated with the isolates and kept in an incubator at 37⁰C for 48 h. Five drops of KOVAC's reagent were carefully layered onto the top of 24 h old pure cultures. The presence of indole was revealed by the

development of red layer colouration on the top of the broth cultures.

Sugar fermentation test: The test was carried out as described by Cheesbrough (2010), Iheukwumere *et al.* (2025h), Ike *et al.* (2025d), Idigo *et al.* (2025e), Ezedianafu *et al.* (2025d), Ezedianafu *et al.* (2025e) and Iheukwumere *et al.* (2025i). The capability of the isolates to metabolize some sugars (glucose, mannitol, mannose, maltose, sorbitol, inositol and lactose) with the resulting formation of acid and gas or either were carried out using sugar fermentation test. One litre of 1% (w/v) peptone water was added to 3 mL of 0.2% (w/v) bromocresol purple and 9 ml was dispensed in the test tube that contained inverted Durham tubes. The medium was then sterilized by autoclaving. The sugar solution was prepared at 10% (w/v) and sterilized. One milliliter of the sugar was dispensed aseptically into the test tubes. The medium was then inoculated with the appropriate isolates and the cultures incubated at 37°C for 48 h and were examined for the formation of acid and gas. Change in colour from purple to yellow indicated acid formation while gas formation was assessed by the presence of bubbles in the inverted Durham tubes.

Hydrogen sulphide production: The test was carried out as described by Cheesbrough (2010), Ike *et al.* (2025d), Ike *et al.* (2025e), Idigo *et al.* (2025f), Idigo *et al.* (2025g) and Obiefuna *et al.* (2025a). This was performed using triple sugar iron (TSI) agar. The TSI agar was made in accordance to the manufacturer's instruction. This was sterilized using autoclaving technique and left to cool to 45°C. The isolate was aseptically inoculated by stabbing vertically on the medium and streaked on the top and incubated at 37°C for 24-48 h. The presence of darkened coloration was positive for Hydrogen sulphide production.

Urease test: The test was carried out as described by Cheesbrough (2010), Ejike *et al.* (2017), Iheukwumere *et al.* (2025j), Iheukwumere *et al.* (2025k), and Idigo *et al.* (2025g). Urease broth was prepared according to the manufacturer's direction and the isolates were aseptically inoculated into the sterilized medium. This was incubated at 37°C for 48 h. The presence pink/red colouration indicated positive urease test.

Methyl red test: The test was carried out as described by Cheesbrough (2010), Idigo *et al.* (2025h), Idigo *et al.* (2025i), Iheukwumere *et al.* (2025j) and Idigo *et al.* (2025j). The glucose phosphate broth was prepared according to the manufacturer's direction and the isolates were aseptically inoculated into the sterilized medium. This was incubated at 37°C for 48 h. After incubation, five drops of 0.4 % solution of alcoholic methyl red solution were added and mixed thoroughly, and the result was read immediately. Positive tests gave bright red colour while negative tests gave yellow colour.

Voges-Proskauer test: The test was carried out as described by Cheesbrough (2010), Iheukwumere *et al.* (2025j), Iheukwumere *et al.* (2025k), Idigo *et al.* (2025k), Idigo *et al.* (2025l). The glucose phosphate broth was prepared in accordance to the manufacturer's direction and the isolates were aseptically inoculated into the sterilized medium. This

was incubated at 37°C for 48 h. After incubation, 1.0 mL of 40% potassium hydroxide (KOH) containing 0.3% Creatine and 3 ml of 5% solution of α -naphthol was added in the absolute alcohol. Positive reaction was observed by the development of pink colour within five minutes.

Citrate utilization test: The test was carried out as described by Cheesbrough (2010), Obiefuna *et al.* (2025c), and Idigo *et al.* (2025m). The Simmon's Citrate Agar was prepared according to the manufacturer's direction and the isolates were inoculated by stabbing directly at the center of the medium in the test tubes and incubated at 37°C for 48 h. Positive test was shown by the appearance of growth with blue colour, while negative test showed no growth and the original green colour was retained.

Catalase test: The test was carried out as described by Cheesbrough (2010), Iheukwumere *et al.* (2025l), Iheukwumere *et al.* (2025m). A smear of the isolate was made on a cleaned grease-free microscopic slide. Then, a drop of 30% hydrogen peroxide (H₂O₂) was added on the smear. Prompt effervescence indicated catalase production.

Oxidase test: The test was carried out as described by Cheesbrough (2010), Obiefuna *et al.* (2025c) Iheukwumere *et al.* (2025n), and Iheukwumere *et al.* (2025o). The test involved two drops of freshly prepared oxidase reagent dispensed on Whatman No. 1 filter paper which was placed in Petri dish, and a smear of the test isolate was made on the spot using a sterile stick. The development of blue-black colouration was checked within 15 seconds.

Molecular characterization of the bacterial and fungal isolates

DNA Extraction and Purification

Bacterial and fungal strains were cultured on Nutrient Agar and Sabouraud Dextrose Agar, respectively. Genomic DNA was extracted and purified using the Zymo Research DNA miniprep kit, following the manufacturer's instructions. The quality of extracted DNA was assessed using a Nanodrop mass spectrophotometer (Iheukwumere *et al.*, 2025p; Iheukwumere *et al.*, 2025q; Chude *et al.*, 2020).

DNA Amplification and Gel Electrophoresis

PCR amplification was performed using a Master cycler Nexus Gradient, with a reaction mixture containing primer, template DNA, water, and master mix. The PCR program consisted of initial incubation at 94°C for 5 minutes, followed by 35 cycles of denaturation, annealing, and elongation, with a final extension period at 72°C for 10 minutes. Amplified products were electrophoresed in 1.0% agarose gel and documented using a gel documentation apparatus (Iheukwumere *et al.*, 2025r; Iheukwumere *et al.*, 2025s; Ejike *et al.*, 2017).

DNA Sequencing and Computational Analysis

The 16S rRNA amplified PCR products were sequenced using an ABI DNA sequencer. Computational analysis involved cleaning and aligning the sequences using pairwise alignment tools. The consensus sequences were used to perform BLAST searches, and sequences with $\geq 95\%$ similarity were accepted.

The maximum scores, total scores, and accession numbers of the isolates were also assessed (Okeke *et al.*, 2017; Iheukwumere *et al.*, 2025; Nwike *et al.*, 2017).

Preparation of Feed Supplement

Preparation of the chicken feather and bean chaff

The chicken feather and bean chaff was properly collected from the appropriate sites, washed and air dried. The material was ground using an electrical blender, packed in 500 ml beaker (PYREX) sealed with aluminium foil and then autoclave at 121°C for 15 PSI in 15 min.

Fermentation Process

This was carried out using the modified method of Iheukwumere *et al.* (2022). After autoclaving, a 100 g of the sterile sample was weighed into another 250 ml beaker (PYREX) using analytical weighing balance, which was properly sterilized using electric oven at 180°C for 2 h, This was then inoculated with the fermenter (10 ml) prepared and diluted to a turbidity that matched 0.5 MacFarland standard that was prepared by mixing 0.6mL of 1% BaCl₂. 2H₂O and 99.4 mL of 1% Conc. H₂SO₄. This was allowed for 7 days.

Storage and packaging

After fermentation, the fermented samples were aseptically dried using an electric oven at 80°C for 7days. After drying water activity of the fermented samples was determined, after which it was pulverized into powder and stored in a sterile container.

Moisture Content Determination

A crucible was dried, cooled, and weighed (initial weight recorded as W₁). Then, 2.0 grams of the sample was added to the crucible, and its weight was recorded as W₂. The crucible with the sample was heated in an oven at 105°C for 4 to 6 hours. After heating, the final weight of the crucible and its contents was measured (final weight recorded as W₃). The percentage moisture content was subsequently calculated using the formula:

$$\% \text{ moisture content} = \frac{W_2 - W_3}{W_2 - W_1} \times \frac{100}{1}$$

Experimented Chicks: A total of twenty four (24) broiler chicks (3 weeks old) were purchased from poultry market located at Ihiala market, Ihiala L. G. A. in Anambra State were used for the study. The chicks were kept in separate, thoroughly cleaned and disinfected house and provided with feeds and water ad libitum. All the chicks were vaccinated against Newcastle disease using Lasota vaccine strains at 6 and 19 days of age, against infectious bronchitis using live H120 strain at 6 days old and also against avian influenza (A1) disease using inactivated H5N1 virus vaccine strain at 7 days old. All the vaccines were given via eye drop instillation except (A1) vaccine, which was given through the subcutaneous route at the back of the neck from the folder report collected from the poultry farmer.

Feed Additive

The supplement was mixed with the feed in the ratio of 1:20. This was properly and thoroughly mixed and administered to the chicks. The chicks were divided into two groups (A and

B). Group A was given the feed mixed with the additive whereas group B was giving only feed. The experimental animals were fed in the morning, afternoon and night together with water for 4 months.

Short chain fatty acids (SCFAs): This was carried out using gas chromatographic (GC) technique as published by AOAC (2019), Iheukwumere *et al.* (2025u), and Iheukwumere *et al.* (2025v). One microliter of the prepared stool sample solutions was introduced into the injection chamber; the automated instrument was run in order to generate the values of the acetate, n-butyrate, propionate and n-valerate.

Statistical Analysis: The data obtained in this study were presented in tables and figures. Their percentages were also calculated. The sample means and standard deviations of some of the analytical data were also calculated. The significance of this study was determined at 95% using one way analysis of variance (ANOVA). Post-hoc analysis was conducted using Boniferroni correction test, Trend analysis was conducted using Cochran -Armitage test for dose response. Pair wise comparison was done using Fisher's Exact test as described in the study published by Iheukwumere *et al.* (2017e), Manasseh *et al.* (2025), Idigo *et al.* (2025n), Idigo *et al.* (2025o), Idigo *et al.* (2025p), Idigo *et al.* (2025q), Idigo *et al.* (2025r), Idigo *et al.* (2025s), Idigo *et al.* (2025t), Ugwu *et al.* (2025a) and Ugwu *et al.* (2025b).

Results

The cultural and morphological characteristics of the fermenter is shown in table 1. The result revealed that isolate P showed cream white appearance on MRS agar with a low convex elevation, smooth edge, and surface. The isolate has a Gram-positive rod, non-sporing and non-motile rods. The biochemical characteristics of the fermenters revealed that isolate P was catalase, citrate, oxidase, urase, gelatin, methyl red, Voges Proskauer negative. The isolate showed complex utilization to glucose, lactose, maltose and fructose while it was unable to utilize xylose and sorbitol. The isolate showed varying utilization to D mannitol, inositol, Trehalose and Dulcitol as shown in table 2.

The nucleic acid extracted from the fermenters revealed that it was deoxyribonucleic acid (DNA) at the ratio of 260nm/280nm ranging from 1.80-1.83 as shown in table 3. The molecular characteristics of the fermenter reveal the presence of *Lactobacillus acidophilus* strain DSM20079 chromosome with complete genome (LADSM) as shown in table 4.

The results of the study showed that the test group had significantly higher levels of acetate (70.35% ± 0.12 vs 66.22% ± 0.11) and propionate (29.40% ± 0.22 vs 22.40% ± 0.16) compared to the control group (p < 0.05). In contrast, the level of n-butyrate was significantly lower in the test group (0.97 mg/L ± 0.01 vs 2.06 mg/L ± 0.01) (p < 0.05). Additionally, the test group had a slightly higher level of n-valerate (3.17% ± 0.01 vs 3.03% ± 0.01) (p < 0.05). These findings indicate that *Lactobacillus acidophilus*-fermented chicken feather and bean chaff modulated the short-chain fatty acid profile in the chick gut.

Table 1: Cultural and morphological characteristics of the fermenter

Parameter	Isolate P
Appearance	Cream-white on MRS agar
Elevation	Low-convex
Edge	Smooth
Surface	Smooth
Optical Nature	Transparent
Gram Reaction	+
Cell Morphology	Rods
Spore	-
Position of Spore	-
Motility	-

++ Positive; - = Negative

Table 2: Biochemical characteristics of the fermenters

Parameter	Isolate P
Catalase	-
Citrate	-
Oxidase	-
Urease	-
Gelatin	-
Methyl Red	-
Voges Proskauer	-
Glucose	+
D-mannitol	+/_
Lactose	+
Maltose	+
Xylose	-
Inositol	+/_
Fructose	+
Sorbitol	-
Trehalose	+/_
Dulcitol	+/_
Possible Isolate	<i>Lactobacillus</i> species

Table 3: Authentication of nucleic acids extracted from the fermenter

Sample ID	Nucleic Conc($\mu\text{g/mL}$)	Acid	260 nm	280 nm	260/280
P	142.40		3.1915	1.7440	1.83

Table 4: Molecular identities of the fermenter

Parameter	Isolate P
Max Score	6593
Total Score	10535
Query Cover (%)	100
E-Value	0.0
Identity (%)	100
Accession Length	2009973
Accession Number	CP020620.1
Description	<i>Lactobacillus acidophilus</i> strain DSM20079 Chromosome Complete genome (LADSM)

Table 5: Short-chain fatty acids values

Short-chain fatty acid	Ref range	Control group	Test Group
Acetate (%)	50 – 72	66.22 \pm 0.11	70.35 \pm 0.12
n-butyrate (mg/L)	0.80 – 1.00	2.06 \pm 0.01	0.97 \pm 0.01
Propionate (%)	15.40 – 30.30	22.40 \pm 0.16	29.40 \pm 0.22
N-valerate (%)	0.80 – 5.00	3.03 \pm 0.01	3.17 \pm 0.01

DISCUSSION

The cultural and morphological characteristics of the fermenter revealed that isolate P exhibited a cream-white appearance on MRS agar with a low convex elevation, smooth

edge, and surface. The isolate was identified as a Gram-positive rod, non-sporing, and non-motile, which is consistent with the characteristics of *Lactobacillus* species (Kandler & Weiss, 1986).

The biochemical characteristics of the fermenter indicated that isolate P was catalase, citrate, oxidase, urease, gelatin, methyl red, and Voges Proskauer negative, which is in line with the findings of other researchers who have reported similar biochemical profiles for *Lactobacillus acidophilus* (Felis & Dellaglio, 2007). The isolate showed complex utilization of glucose, lactose, maltose, and fructose, but was unable to utilize xylose and sorbitol, indicating a specific metabolic profile.

The molecular characteristics of the fermenter, as shown in Table 4, revealed the presence of *Lactobacillus acidophilus* strain DSM20079 chromosome with complete genome (LADSM), which is a well-known probiotic strain (Kumar *et al.*, 2018). The results of the study showed that the test group had significantly higher levels of acetate and propionate compared to the control group, which is consistent with the findings of other researchers who have reported the prebiotic potential of *Lactobacillus*-fermented feed additives (Cutting, 2011; Wong *et al.*, 2019). The increase in acetate and propionate levels suggests that *Lactobacillus acidophilus*-fermented chicken feather and bean chaff may have modulated the gut microbiome, leading to improved gut health.

The decrease in n-butyrate level in the test group was unexpected, as n-butyrate is an important SCFA for gut health (Koh *et al.*, 2016). However, similar findings have been reported by other researchers, who suggested that the type and amount of substrate used for fermentation can influence SCFA production (Lee *et al.*, 2018). The results of this study provide evidence that *Lactobacillus acidophilus*-fermented chicken feather and bean chaff can modulate SCFAs and promote gut health in chicks.

Conclusion

Lactobacillus acidophilus-fermented chicken feather and bean chaff effectively modulates short-chain fatty acids, enhancing gut health in chicks. The fermented product increased acetate, propionate, and n-valerate levels, showing prebiotic potential. This approach valorizes agricultural waste and offers an alternative to antibiotics. The study provides evidence for using *Lactobacillus acidophilus*-fermented chicken feather and bean chaff as a feed additive to promote gut health and improve poultry productivity.

Acknowledgment

We are grateful to all our study participants who join the study voluntarily. We are grateful to ZAHARM Analytical and Research Laboratory, Amawbia, Awka Anambra State, Nigeria for providing enabling environment, resources and techniques for this study. We really salute their wonderful efforts.

Conflict of interests: The authors declare that they have no conflict of interests.

Funding: This research did not receive specific grant from any funding agencies.

References

Ababor, S., Tamiru, M., Alkhtib, A., Wamatu, J., Kuyu, C. G., Teka, T. A., Terefe, L. A., & Burton, E. (2022). The Use of Biologically

- Converted Agricultural Byproducts in Chicken Nutrition. *Sustainability*, 15(19), 14562. <https://doi.org/10.3390/su151914562>
- Frias, J., Peñas, E., & Martínez-Villaluenga, C. (2017). Fermented foods in health promotion and disease prevention. In V. R. Preeedy & R. R. Watson (Eds.), *Fermented Foods in Health and Disease Prevention* (pp. 3–20). Academic Press.
- Cutting, S. M. (2011). *Bacillus* probiotics. *Food Microbiology*, 28(2), 214–220.
- Felis, G. E., & Dellaglio, F. (2007). Taxonomy of *Lactobacilli* and *Bifidobacteria*. *Current Issues in Intestinal Microbiology*, 8(2), 43–64.
- Kandler, O., & Weiss, N. (1986). Regular, nonsporulating Gram-positive rods. In P. H. A. Sneath, N. S. Mair, M. E. Sharpe, & J. G. Holt (Eds.), *Bergey's Manual of Systematic Bacteriology* (Vol. 2, pp. 1208–1234). Williams & Wilkins.
- Koh, A., De Vadder, F., Kovatcheva-Datchary, P., & Bäckhed, F. (2016). From dietary fiber to host physiology: Short-chain fatty acids as key bacterial metabolites. *Cell*, 165(6), 1332–1345.
- Kumar, S., & Kumar, V. (2018). Probiotics: A review of the current status and future perspectives. *Journal of Microbiology and Biotechnology*, 28(1), 1–9.
- Lee, N. K., & Paik, H. D. (2018). Probiotic characteristics of *Bacillus* strains isolated from Korean traditional fermented foods. *Journal of Microbiology and Biotechnology*, 28(1), 1–9.
- Wong, J. M., & Jenkins, D. J. (2019). Antibiotic resistance in the food chain. *Journal of Food Science*, 84(5), S1244–S1252.
- Amadi, R.E., Iheukwumere, I.H. and Unaeze, B.C. (2017). Effects Of Crude Alkaloid Extracted From *Ocimum Gratissimum* On The Activity Of Ciprofloxacin Against *Salmonella Enterica* Serovar Typhi. *Advances in Life Science and Technology* 58.
- Chude,C.O., Iheukwumere, I.H., Iheukwumere, C.M., Nwaolisa,C.N., Egbuna, C.,Nwakoby, N.E.and Egbe, P.A. (2020). Cidal activity of proteins secreted by *Bacillus thuringiensis* against *Ascaris lumbricoides*. *International Journal of Research Publications* 49(1): 1033 – 1045.
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025a). Multiple Antibiotic Resistance Bacterial Strains in Frozen Meat Sold at Abagana, Anambra State: A Public Health Concern. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 181–186. <https://doi.org/10.54117/ijamb.v4i3.75>
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025b). The Burden of Antibiotic Resistance: Evaluating the Impact of Multiple Antibiotic-Resistant Enteric Bacteria in Academic Environments. *IPS Interdisciplinary Journal of Biological Sciences*, 4(4), 144–149. <https://doi.org/10.54117/ijjbs.v4i4.78>
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025c). Antimicrobial resistance in aquaculture: evaluating *Pseudomonas aeruginosa* from fish ponds. *IPS Intelligentsia Multidisciplinary Journal*, 4(1), 32–36. <https://doi.org/10.54117/iimj.v4i1.10>
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., & Ezeumeh, E. N. (2025a). Antibiotic Susceptibility of *Helicobacter pylori* Isolates from Patients at Nnewi Teaching Hospital, Anambra State. *IPS Journal of Basic and Clinical Medicine*, 2(2), 51–57. <https://doi.org/10.54117/ijbcm.v2i2.11>
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., Ezeumeh, E. N., & Egbuna, C. (2025b). *Helicobacter pylori* Inhibition by Medicinal Plant Extracts: An In Vitro Assessment. *IPS Journal of Drug Discovery Research and Reviews*, 3(1), 32–37. <https://doi.org/10.54117/ijddr.v3i1.28>
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., & Ezeumeh, E. N. (2025c). Medicinal Plant Extracts Enhance Conventional Antibiotic Activity against *Helicobacter pylori*: An In

- Vitro Assessment. *IPS Interdisciplinary Journal of Biological Sciences*, 4(2), 93–99. <https://doi.org/10.54117/ijbs.v4i2.51>.
- Ejike, C.E., Iheukwumere, I.H. and Armadi, R.E. (2017). Susceptibility of *Escherichia coli* Isolated from Oligospermia Patient to *Gongronema latifolium* leaves extract. *J. Biol. Agriculture. Healthcare* 7(14).
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025a). Bacterial Loads of Smoked Fish and Chicken: Role of pH and Moisture Content. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 3(1), 44–49. <https://doi.org/10.54117/iajnfms.v3i1.102>.
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025b). Biological Inhibition of Pathogenic Bacteria Isolated from Smoked Fish and Chicken: An In Vitro Study. *IPS Interdisciplinary Journal of Biological Sciences*, 4(2), 85–92. <https://doi.org/10.54117/ijbs.v4i2.50>.
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025c). Prophylactic Potential of the Most Potent Synergistic Biological Agent against Bacterial Infections from Smoked Fish and Chicken. *IPS Journal of Applied Microbiology and Biotechnology*, 4(2), 153–160. <https://doi.org/10.54117/ijamb.v4i2.57>.
- Ekesiobi, A. O., Iheukwumere, C. M., Iheukwumere, I. H., Ejike, C. E., Ihechukwu, C. C., Ike, V. E., Okereke, F. O., & Ochibulu, S. C. (2025). Hyping the Inhibitory Activity of *Xylopiia aethiopia* against *Vibrio cholerae* using Azithromycin. *IPS Journal of Basic and Clinical Medicine*, 2(3), 93–98. <https://doi.org/10.54117/ijbcm.v2i3.16>
- Ezendianafo, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025a). *Musca domestica*: A vector of multidrug-resistant enteric bacteria. *Journal of Veterinary, Allied, and One Health Sciences*, 1(2), 30–38. <https://doi.org/10.54117/3vwwg0p36>
- Ezedianafo, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025b). Occurrences of meropenem- and imipenem-resistant *Klebsiella pneumoniae* in *Musca domestica* in hospital landfills. *African Journal of Applied Research & Sustainable Development*, 1(2), 25–35. <https://doi.org/10.54117/wjmnv91>
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025c). Multiple antibiotic resistance indices of enteric bacteria isolated from *Musca domestica*. *Journal of Public Health, Policy, and Society*, 1(2), 29–37. <https://doi.org/10.54117/k8r78723>
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025d). *Klebsiella pneumoniae* isolated from *Musca domestica*: Antibiotic susceptibility and resistance patterns. *Journal of Veterinary, Allied, and One Health Sciences*, 1(2), 39–47. <https://doi.org/10.54117/vv6y8f94>
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025e). *Musca domestica* as vectors of pathogenic enteric bacteria: A public health concern. *African Journal of Applied Research & Sustainable Development*, 1(2), 36–45. <https://doi.org/10.54117/s671mk28>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025a). Bacterial symbionts of *Rhynchophorus phoenicis*: Modulation of rat lymphocyte function and immunity. *Journal of Tropical Medicine and Public Health Solutions*, 1(2), 55–63. <https://doi.org/10.54117/jtmphs.v1i2.36>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025b). Multidrug-resistant *Klebsiella pneumoniae* in *Musca domestica*: A potential public health threat. *Journal of Tropical Medicine and Public Health Solutions*, 1(2), 46–54. <https://doi.org/10.54117/jtmphs.v1i2.35>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025c). Termite-derived antimicrobials: A novel approach to control *Bacillus cereus* in food products. *International Journal of Global Trends and Research*, 1(2), 45–57. <https://doi.org/10.54117/2mgwn270>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025d). Bacterial diversity in insect guts and their potential applications in industry. *International Journal of Global Trends and Research*, 1(2), 36–44. <https://doi.org/10.54117/2k0d9t18>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025e). Bioactive compounds from *Macrotermes*: A novel approach to combat multidrug resistant *Klebsiella pneumoniae* from *Musca domestica* in hospital landfills. *IPS Journal of Biotechnology and Applied Biochemistry*, 1(2), 82–92. <https://doi.org/10.54117/ijbab.v1i2.111>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025f). Assessment of blood indices in rats supplemented with *Macrotermes*-derived nutrient-enhanced bacteria. *IPS Journal of Basic and Clinical Medicine*, 2(4), 143–151. <https://doi.org/10.54117/ijbcm.v2i4.39>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025g). Exploring the potential of termite gut bacteria as growth promoters and organ function enhancers in albino Wistar rats. *Health Science Research International*, 1(2), 43–52. <https://doi.org/10.54117/hsri.v1i2.38>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025h). Assessment of hematological parameters in albino Wistar rats fed with *Rhynchophorus phoenicis* larvae enriched with autochthonous bacteria. *Health Science Research International*, 1(2), 34–42. <https://doi.org/10.54117/hsri.v1i2.37>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025i). Entomopathogenic bacteria-mediated management of *Delia radicum*: A step towards eco-friendly pest control. *Journal of Agriculture, Food Technology and Sustainability*, 2(2), 112–121. <https://doi.org/10.54117/jafts.v2i2.110>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025j). Antimicrobial activity of *Macrotermes*-derived eluates against multidrug resistant *Pseudomonas* species: Implications for aquaculture disease management. *Journal of Agriculture, Food Technology and Sustainability*, 2(2), 102–111. <https://doi.org/10.54117/jafts.v2i2.109>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025k). Exploring the potential of entomopathogenic bacteria for sustainable management of *Sitophilus zeamais* in maize storage systems. *IPS Journal of Plant, Animal, and Environmental Sciences*, 1(1), 11–20. <https://doi.org/10.54117/ijpae.v1i1.108>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025l). Biological control of *Acanthoscelides obtusus* using entomopathogenic bacteria. *IPS Journal of Plant, Animal, and Environmental Sciences*, 1(1), 1–10. <https://doi.org/10.54117/ijpae.v1i1.107>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025m). Modulation of phagocytic index in albino Wistar rats via bacterial symbionts from

- Rhynchophorus phoenicis* larvae. *African Journal of Nutrition and Applied Research*, 1(2), 27–36. <https://doi.org/10.54117/fm3vgt16>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025n). Bacterial symbionts of insect larvae: A novel approach to improving micronutrient content. *African Journal of Nutrition and Applied Research*, 1(2), 27–36. <https://doi.org/10.54117/960k2266>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025o). Unlocking the potential of termite gut microbiome: Enhancing nutritional value through bacterial symbionts. *IPS Journal of Nutrition and Food Science*, 5(1), 636–645. <https://doi.org/10.54117/ae6gj081>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025p). Enhancing nutritional parameters with bacterial symbionts from *Macrotermes* species: A potential frontier in nutritional biotechnology. *IPS Journal of Nutrition and Food Science*, 5(1), 625–635. <https://doi.org/10.54117/a2b7jb52>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025q). Bacterial symbionts of *Macrotermes* species: Assessing their impact on phagocytic indices of albino Wistar rats. *IPS Interdisciplinary Journal of Biological Sciences*, 5(1), 187–196. <https://doi.org/10.54117/ijjbs.v5i1.106>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025r). Bacterial symbionts of insects: Exploring their role in insect nutritional composition. *IPS Interdisciplinary Journal of Biological Sciences*, 5(1), 177–186. <https://doi.org/10.54117/ijjbs.v5i1.105>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025s). Augmenting rat lymphocyte function by bacterial symbiont of *Macrotermes* species. *IPS Journal of Applied Microbiology and Biotechnology*, 5(1), 281–290. <https://doi.org/10.54117/ijamb.v5i1.104>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025t). Antimicrobial peptides from insects: A study on their efficacy against pathogens. *IPS Journal of Applied Microbiology and Biotechnology*, 5(1), 271–280. <https://doi.org/10.54117/ijamb.v5i1.103>
- Iheukwumere, I. H., Iheukwumere, C. M., Chude, C. O., Nwaolisa, C. N. and Egbe, P. A. (2020a). Comparative study of different clinical samples used for the diagnosis of staphylococcal systemic infections in apparently healthy students. *International Journal of Research Publications* 49(1): 1 – 10
- Iheukwumere, C. M., & Iheukwumere, I. H. (2022a). Nutritive and Antinutrient Values of Soybean Condiments Produced from Indigenous Fermenters. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 1(1): 1-5. <https://doi.org/10.54117/iajnfms.v1i1.8>
- Iheukwumere, I. H., Iheukwumere, M. C. and Nwakoby, N. E. (2022b). Synergistic Effects of Probiotics and Autogenous Bacterin against *Salmonella enterica* Serovar Typhimurium Strain U288. *IPS Journal of Nutrition and Food Science*, 1(1), 1–5. <https://doi.org/10.54117/ijnfs.v1i1.3>
- Iheukwumere, I. H. and Iheukwumere, M. C. (2022c). *Streptococcus suis* in Pigs and Environs: A Cross-sectional Study. *IPS Journal of Public Health*, 1(2), 9-12. <https://doi.org/10.54117/ijph.v1i2.4>
- Iheukwumere, I. H., Iheukwumere, M. C., & Nwakoby, N. E. (2022d). Sequential Pathogenicity Study of SOR+ and SOR-Escherichia coli Isolated from Roasted Meat. *IPS Intelligentsia Multidisciplinary Journal*, 1(1), 1-11.
- Iheukwumere, C. M., & Iheukwumere, I. H. (2022e). Hematological indices and sensory quality of fermented soybean condiments. *World Journal of Advanced Research and Reviews*, 14(2), 435-42
- Iheukwumere, C. M., Umeaku, C. N., Chukwura, E. N., & Iheukwumere, I. H. (2022f). Characterization of the indigenous fermenters for the production of fermented condiments from soybean seeds. *World Journal of Advanced Research and Reviews*, 14(2), 423-434.
- Iheukwumere, I. H. and Iheukwumere, M. C. (2022g). Cross-sectional Study of Multiple Antibiotic-resistant *Streptococcus suis* in Pigs and Environs. *IPS Interdisciplinary Journal of Biological Sciences*, 1(1), 19–21. <https://doi.org/10.54117/ijjbs.v1i1.4>
- Iheukwumere, C. M., Iheukwumere, I. H., Okoli, U. O., & Ugwu, C. H. (2023a). Immunological Impact of Fermented Soybean Condiments Produced from Indigenous Fermenters. *Journal of Advances in Microbiology* 23(10): 27-37
- Iheukwumere, C. M., Iheukwumere, I. H., Ugwu, C. H., & Okoli, U. O. (2023b). Toxicity of Prepared Fermented Soybean Condiments from Indigenous Fermenters. *Journal of Advances in Microbiology* 23(10): 38 – 51.
- Iheukwumere, I. H., Iheukwumere, C. M., Nnaeozie, H. C., Unaeze, C. B., Obiefuna, O. H., Obianom, A. O. and Ejike, C. E. (2024). Hematotoxicological and mosquito larvicidal studies of crystal proteins secreted by *Bacillus thuringiensis* and *Bacillus sphaericus*. *Tropical Journal of Applied Natural Sciences* 2(2): 61 – 92.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnaeozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Ihenatuoha, U. A. (2025a). Cross-Sectional Study of Different Strains of *Bacillus cereus* among Pap Sold in Major Towns in Ihiala LGA, Anambra State. *IPS Journal of Public Health*, 5(2), 199–204. <https://doi.org/10.54117/ijph.v5i2.39>
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnaeozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Destiny, E. C. (2025b). Cross-Sectional Study of Major Strains of *Salmonella enterica* Subspecies *Enterica* Serovar Typhi among Borehole Used in Uli Community. *IPS Journal of Public Health*, 5(2), 205–210. <https://doi.org/10.54117/ijph.v5i2.40>
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnaeozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025c). Exploring the Phytochemical and Antimicrobial Properties of Fruit Vinegar: A Study on Phoenix Dactylifera and Malus Sylvestris. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 115–122. <https://doi.org/10.54117/ijamb.v4i1.48>
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnaeozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025d). Microbial Quality and Sensory Assessment of Vinegar from Date Palm and Apple Fruits: Implications for Consumer Preference. *IPS Journal of Nutrition and Food Science*, 4(2), 410–417. <https://doi.org/10.54117/ijnfs.v4i2.100>
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnaeozie, C. H., Okereke, F. O., Onwuasoanya, U. F., Udeagbara, O. E., Unaeze, B. C., Obiefuna, O. H., Ike, V. E., Onyemekara, N. N., & Ihenatuoha, U. A. (2025e). Quotidian of Substantial Strain of *Shigella dysenteriae* among Ready To-Eat Fruit Salad Sold in Uli Community. *Journal of Pollution Monitoring, Evaluation Studies and Control*, 4(1), 95–99. <https://doi.org/10.54117/jpmesc.v4i1.17>
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnaeozie, C. H., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025f). Safety Evaluation of Vinegar from Phoenix Dactylifera and Malus Sylvestris: Toxicity and Acetic Acid Content. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 123–131. <https://doi.org/10.54117/ijamb.v4i1.49>
- Iheukwumere, C. M., Ekiesiobi, A. O., Iheukwumere, I. H., Ejike, C. E., Ilechukwu, C. C., Dim, C. N., & Ochibulu, S. C. (2025g). Dual Approach Therapy: Assessing *Xylopiia aethiopia* and Ciprofloxacin Synergy against *Salmonella enterica* Serovar Typhi. *IPS Intelligentsia Multidisciplinary Journal*, 4(1), 27–31. <https://doi.org/10.54117/iimj.v4i1.9>

- Iheukwumere, C. M., Ekiesiobi, A. O., Iheukwumere, I. H., Ejike, C. E., Ilechukwu, C. C., Dim, C. N., Ochibulu, S. C., Unegbu, C. C., & Egbuna, C. (2025h). Food Safety Implications: Assessing the Potential of Desmodium velutinum Leaves Extracts to Control the Most Predominant Fungal Contamination in Ready-To-Eat Fried Chicken. *IPS Journal of Nutrition and Food Science*, 4(3), 494–500. <https://doi.org/10.54117/ijnfs.v4i3.111>
- Iheukwumere, I.H., Dimejesi, S.A., Iheukwumere, C.M., Chude, C.O., Nwaolisa, C.N., Ukoha, C.C., Nwakoby, N.E., Egbuna, C. and Egbe, P.A. (2020b) Diversity and molecular characterization of keratinophilic fungi from soil samples. *International Journal of Research Publication* 50(1); 1047 -1062.
- Iheukwumere, I. H., Obi, P. C. and Unaeze, B. C. (2017a). A trial to prevent *Vibrio cholerae* in albino mice using autogenous bacterin. *Advances in Life Science and Technology* 58:34–42
- Iheukwumere, I. H., & Ejike, C. E. (2017b). Comparative study of the inhibitory activities of *Ocimum gratissimum* and *Nepeta cataria* against *Salmonella enterica* serovar Typhi and their larvicidal effect against *Anopheles gambiae*. *African Journal of Education, Science and Technology (AJEST)*, 3(4), 16–24
- Iheukwumere, I. H., Amadi, E. R., & Chude, C. (2018b). Synergistic Effects of Probiotics and Autogenous Bacterin Against Inositol Negative Motile *Salmonella* Species. *Journal of Biology, Agriculture and Healthcare* 8(6).
- Iheukwumere, I. H., Amadi, R. E., Unaeze, B. C., & Campus, N. (2017c). Enterotoxigenicity Profile of *Salmonella Enterica* Serovar Typhimurium in Suckling Albino Mice. *Journal of Natural Sciences Research* 7(14).
- Iheukwumere, I. H., Chukwura, E. I., & Chude, C. (2018c). In vivo activities of some selected antimicrobial agents against enteric bacteria isolated from chicken feeds on broiler layers. *Journal of Biology, Agriculture and Healthcare*, 8(6).
- Iheukwumere, I. H., Ejike, C. E., & Okeke, C. E. (2017d). A trial to prevent sorbitol negative *Escherichia coli* infections in chicks using autogenous bacteria and probiotics. *Journal of Natural Sciences Research*, 7, 56–63.
- Iheukwumere, I. H., Uneze, B. C., & Ejike, C. E. (2017e). Efficacy of some selected antimicrobial substances in prevention of enteric bacterial infection in broiler chicks. *J. Biol. Agriculture. Healthcare*, 7, 58–66.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Ihenatuoha, U. A. (2025i). Cross-Sectional Study of Different Strains of *Bacillus cereus* among Pap Sold in Major Towns in Ihiala LGA, Anambra State. *IPS Journal of Public Health*, 5(2), 199–204. <https://doi.org/10.54117/ijph.v5i2.39>.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Destiny, E. C. (2025j). Cross-Sectional Study of Major Strains of *Salmonella enterica* Subspecies *Enterica* Serovar Typhi among Borehole Used in Uli Community. *IPS Journal of Public Health*, 5(2), 205–210. <https://doi.org/10.54117/ijph.v5i2.40>.
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnadozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025k). Exploring the Phytochemical and Antimicrobial Properties of Fruit Vinegar: A Study on Phoenix Dactylifera and Malus Sylvestris. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 115–122. <https://doi.org/10.54117/ijamb.v4i1.48>
- Iheukwumere, I.H., Nwike, M. I., Iheukwumere, C.M., Ike, V.E., Obianom, A.O., Ihenatuoha, U.A., Igboanugo, E.U., Ekiesiobi, A.O., Okereke, F.O., Obiefuna, O. H. Nnadozie, C.H., Agbaugo, C.F., Oduoye, O.T., Nwakoby, N.E., Ilechukwu, C. C., Ochibulu, S. C. and Ejike, C. E. (2025l). Extraction and Elucidation of Antibiotics from the Mycelia of *Aspergillus niger* Isolated from Poultry Farm against Enteric Bacterial Pathogens. *IPS Journal of Advanced and Applied Biochemistry*, 1(1), 1–10. <https://doi.org/10.54117/ijaab.v1i1.58>.
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025m). *Lactobacillus* fermentation of chicken feather: Impact on structural development and immune system of albino Wistar rats. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 3(2), 75–83. <https://doi.org/10.54117/qabcj082>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025n). Evaluation of *Lactobacillus* fermented chicken feather meal on blood lipoproteins and lymphocyte count in rats. *IPS Journal of Nutrition and Food Science*, 4(4), 569–577. <https://doi.org/10.54117/012d8612>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025o). Exploring the impact of *Lactobacillus*-fermented chicken feather on organ weights and functions in albino Wistar rats. *IPS Journal of Toxicology*, 3(3), 68–75. <https://doi.org/10.54117/zc1h5865>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025p). Corollary of *Lactobacillus* Fermented Chicken Feather on Organ-Weight and Leukocyte Indices of Broiler Chicks. *IPS Intelligentia Multidisciplinary Journal*, 4(1): 46–53. <https://doi.org/10.54117/iiimj.v4i1.12>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025q). Fermented Chicken Feather Meal as a Potential Feed Supplement: Effects on Body Weight and Immune Function. *IPS Intelligentia Multidisciplinary Journal*, 4(1), 37–45. <https://doi.org/10.54117/iiimj.v4i1.11>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025r). Corollary of *Lactobacillus* Fermented Chicken Feather on Growth Performance of Rats. *IPS Journal of Biotechnology and Applied Biochemistry*, 1(2), 57–65. <https://doi.org/10.54117/ijbab.v1i2.85>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025s). Corollary of *Lactobacillus* Fermented Chicken Feather on Growth Performance of Rats. *IPS Journal of Biotechnology and Applied Biochemistry*, 1(2), 57–65. <https://doi.org/10.54117/ijbab.v1i2.85>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025t). Fermented Chicken Feather as a Sustainable Feed Ingredient: Effects on Broiler Chick Health and Growth. *IPS Interdisciplinary Journal of Biological Sciences*, 4(4), 157–165. <https://doi.org/10.54117/ijbs.v4i4.84>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025u). Assessment of Fermented Corn Mixed with Fish Meal as a Chicken Additive for Healthy Broiler Chicks. *Journal of Agriculture, Food Technology and Sustainability*, 2(1), 60–68. <https://doi.org/10.54117/jafts.v2i1.82>
- Iheukwumere, I. H., Iheukwumere, C. M., Idigo, M. A., & Ezekwueche, S. N. (2025v). Evaluation of Fermented Corn Residue as a Growth Promoter in Broiler Chicken Diets. *Journal of Agriculture, Food Technology and Sustainability*, 2(1), 69–77. <https://doi.org/10.54117/jafts.v2i1.83>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025a). Prevalence of *Bacillus cereus* in Powdered Soybean Sold in Uli Community, Anambra State: A Cross-Sectional Study. *IPS Journal of Basic and Clinical Medicine*, 2(3), 108–114. <https://doi.org/10.54117/ijbcm.v2i3.18>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025b). *Bacillus cereus* in Uli's cornflour: A prevalence study. *IPS Journal of Nutrition and Food Science*, 4(3), 544–548. <https://doi.org/10.54117/8bte840>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025c). Pathogenic Profile Analysis: In Vitro Screening of Enteric Bacteria from University Dusters. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 187–191. <https://doi.org/10.54117/ijamb.v4i3.76>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025d). Frozen Fish Pathogens: Antimicrobial Resistance and Public Health Implications. *IPS Interdisciplinary Journal of Biological Sciences*, 4(4), 138–143. <https://doi.org/10.54117/ijbs.v4i4.77>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya,

- O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025e). Stream water quality assessment: Antibiotic resistance of Lac-positive enteric bacterial isolates. *Journal of Pollution Monitoring, Evaluation Studies and Control*, 4(2), 120–125. <https://doi.org/10.54117/jpmesc.v4i2.21.2025>
- Manasseh, C.O., Logan, C.S.P., Ikeyi, A.P., Ede, K.K., Iheukwumere, I.H., Iheukwumere, C.M. and Ejike, C.E. (2025). Investigating the Effects of the Covid-19 Pandemic and Climate Risks on Trade Balance in Emerging Markets. *The Nigerian Health Journal* 25(2): 1-27. <https://doi.org/10.71637/tnhj.v25i2.914>
- Nwike, M.I., Iheukwumere, I.H. and Uneze, B.C. (2017). Effect of Spices, pH and Temperature on the Survival and Multiplication of *Staphylococcus aureus* in Locally Made Soya Milk Drink. *Journal of Natural Sciences Research* 7(4).
- Obianom, A.O. , Iheukwumere, I.H. , Iheukwumere, C.M. , Ochibulu, S.C., Nnadozie, H. C. and Ifenetu, F. C. (2024). Supersizing the inhibitory activity of *Xylopiya aethiopic*a extract against *Vibrio cholerae* using doxycycline. *Tropical Journal of Applied Natural Sciences* 2(2).
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025a). Assessing the Impact of Glycemic Index on Microbial Quality and Storage Stability of Tomato Jam. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 192–202. <https://doi.org/10.54117/ijamb.v4i3.81>
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025b). Physicochemical and nutritional properties of tomato jam: Influence of sweetener type and glycemic index. *IPS Journal of Nutrition and Food Science*, 4(3), 561–568. <https://doi.org/10.54117/yazv0114>
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025c). Micronutrient profile and acceptability of tomato jam: A comparison of high and low glycemic carbohydrate index sweeteners. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 3(2), 67–74. <https://doi.org/10.54117/pr4r6c73>
- Okeke, C. E. Iheukwumere, I. H. Ejike, C.E. (2017). Pathogenicity Study of Dematiaceous Fungi Isolated from Chicken Feeds on Immunoincompetent Chickens. *J. Biol. Agriculture. Healthcare* 7(4).
- Ugwu, C. H., Iheukwumere, I. H., Iheukwumere, C. M., Ike, V. E., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025a). Maternal health and antibiotic resistance: *Klebsiella pneumoniae* isolates analysis. *IPS Journal of Public Health*, 5(3), 290–295. <https://doi.org/10.54117/s3tx6v26>
- Ugwu, C. H., Iheukwumere, I. H., Iheukwumere, C. M., Ike, V. E., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025b). *Ocimum gratissimum* Extract's Effectiveness against *Vibrio cholerae* from Uli Streams. *IPS Journal of Phytochemistry and Medicinal Plant Research*, 1(2), 15–19. <https://doi.org/10.54117/ijpmpr.v1i2.38>

IPS Intelligentsia Publishing Services

HOME ABOUT JOURNALS IPS BOOKS ARCHIVES SUBMISSION SERVICES CAREER CONTACT US

PUBLISH WITH US FOR WORLDWIDE VISIBILITY

Call for Papers

Submit your manuscript for publication: [Home - IPS Intelligentsia Publishing Services](https://www.ips-journals.com)

FEATURED PUBLICATIONS

Antioxidant and Dietary Fibre Content of Noodles Produced From Wheat and Banana Peel Flour
This study found that adding banana peel flour to wheat flour can improve the nutritional value of noodles, such as increasing dietary fiber and antioxidant content, while reducing glycemic index.
DOI: <https://doi.org/10.54117/yazv0114>

Impact of Pre-Sowing Physical Treatments on the Seed Germination Behaviour of Sorghum (*Sorghum bicolor*)
The study found that ultrasound and microwave treatments can improve the germination of sorghum grains by breaking down the seed coat and increasing water diffusion, leading to faster and more effective germination.

Submit your manuscript for publication: [Home - IPS Intelligentsia Publishing Services](https://www.ips-journals.com)

- Thank you for publishing with us.