



Fermented Corn Chaff as a Sustainable Feed Additive for Broiler Chick Production

Abba, O.¹, Iheukwumere, I. H.², Iheukwumere, C. M.³, Ike, V. E.⁴, Ezendianefo, J. N.⁵ and Okongwu, D. J.⁶

¹Department of Microbiology, Federal University of Gusau, Zamfara State.

²Department of Microbiology, Faculty of Natural Sciences, Chukwuemeka Odumegwu Ojukwu University, Uli, Campus, Anambra State, Nigeria.



³Department of Applied Microbiology and Brewing, Faculty of Biosciences, Nnamdi Azikiwe University, Awka, Nigeria.

⁴Department of Microbiology, University of Agriculture and Environmental Sciences, Umuagwo, Imo State, Nigeria.

⁵Department of Microbiology, Tansian University, Umunya, Anambra State, Nigeria.

⁶Department of Chemistry, Nwafor Orizu College of Education, Nsugbe.

*Corresponding author email: oluchiipan@fufusau.edu.ng / ik.iheukwumere@coou.edu.ng

Abstract	Article History
<p>The increasing demand for poultry products necessitates sustainable feed additives. Corn chaff, an agricultural waste, is underutilized. While fermentation enhances nutritional value, its impact on broiler chick production remains unclear, leaving a gap in understanding its effects on growth performance and gut health, creating a need to investigate its potential as a sustainable feed additive.. This study evaluated the effect of fermented corn chaff as a chicken additive for healthy broiler chicks. The fermenter, identified as <i>Lactobacillus acidophilus</i> strain DSM20079 (LADSM), was characterized culturally, morphologically, and biochemically. The study showed significant increases ($p < 0.05$) in body weights of test chicks from week 1 (186 g) to week 6 (1824 g) compared to controls (1442 g). Organ weights were similar between groups ($p > 0.05$). Feed conversion ratio (FCR) was lower in test group from week 3-6 ($p < 0.05$). Hematological indices revealed increased WBC (13.68 vs $12.88 \times 10^9/L$), RBC (7.89 vs $7.37 \times 10^{12}/L$), and lymphocytes (73.30% vs 50.70%) in test group ($p < 0.05$). The study concludes that fermented corn chaff exhibits growth-promoting activity and can be used as an additive for healthy broiler chicks.</p> <p>Keywords: Fermented corn chaff, <i>Lactobacillus acidophilus</i>, Broiler chicks, Growth performance, Feed additive</p>	<p>Received: 05 Jan 2026 Accepted: 13 Feb 2026 Published: 17 Feb 2026</p>  <p>Scan QR code to view*</p> <p>License: CC BY 4.0*</p>  <p>Open Access article.</p>
<p>How to cite this paper: Abba, O., Iheukwumere, I. H., Iheukwumere, C. M., Ike, V. E., Ezendianefo, J. N., & Okongwu, D. J. (2026). Fermented Corn Chaff as a Sustainable Feed Additive for Broiler Chick Production. <i>IPS Journal of Applied Microbiology and Biotechnology</i>, 6(1), 317–327. https://doi.org/10.54117/ijamb.v6i1.132</p>	

Introduction

Food production is an indispensable activity that sustains mankind in the universe (Ditto *et al.*, 2018). Production of chicken is highly essential in society due to its role as a food source for humans and economic purposes (Khan and Iqbal, 2016; Iheukwumere *et al.*, 2025a; Dim *et al.*, 2025a). Chickens are domestic animals that need a high level of care, especially when large production is practiced, unlike small-scale producers, who allow chickens to roam in the environment to source food, though there could be feed provision in low quantity (Papatsiros *et al.*, 2013).

It is worthy to note that the type of feed provided to broiler chicks determine their growth rate and overall health of the birds (Ditto *et al.*, 2018; Iheukwumere *et al.*, 2022a; and Nwike *et al.*, 2017). Broiler chicken are specifically raised for

meat production and administering adequate feed in good proportion is paramount (Ditto *et al.*, 2018).

In the society, the use of chemical compounds to enhance the growth of broiler chicks is alarming (Correa-Oliveira *et al.*, 2016; Ditto *et al.*, 2018; Ekechukwu *et al.*, 2025a; Obianom *et al.*, 2024; Dim *et al.*, 2025b). These chemical compounds facilitate quick growth but could pose health risk to consumers via bio-magnification. Most of the diseases that are ravaging the health of individuals had been attributed to chemicals found in processed foods and poultry as well (Awad *et al.*, 2017; Iheukwumere *et al.*, 2025b, Dim *et al.* 2025c).

Research has revealed that natural additives and feeds are capable of providing enormous health benefits to both the broiler chicks and the consumers (Mcknight *et al.*, 2020; Sun

et al., 2020). Natural wastes from crops such as corn chaff and beans chaff contain fibre, which when fermented by certain microorganisms such *Bacillus* species, *Lactobacillus* species, and *Saccharomyces cerevisiae* produces essential byproducts (Amadi *et al.*, 2017; Ejike *et al.*, 2017; Latorre *et al.*, 2018). Microbial fermentation of fibre in crops produces essential metabolites that confer health benefits to broiler chicken such as antimicrobial activity, regulation of healthy intestinal mucosa, growth of intestinal cells, and modulation of immune response (Gharib-Naaseri *et al.*, 2019) and short chain fatty acid production during fermentation contributes immensely in the aforementioned benefits (Onrust *et al.*, 2018).

Several studies are available on the role of fermentative products on the growth performance in broiler chicks such as Onrust *et al.* (2018), Adhikari *et al.* (2020), and Aljumaah *et al.* (2020) but few studies are available on the assessment of fermented corn chaff as chicken additive for healthy broiler chicks. Hence, this study aims to assess fermented corn chaff as chicken additive for healthy broiler chicks.

Materials and Methods

Isolation of the Test Sample

The media used for this isolation was de Man Rogosa and Sharpe broth (MRS) (BIOTECH). A 1.0 ml of fermented yoghurt (Aqua yoghurt) and banana extract were aseptically introduced into sterile Petri dishes (90 mm x 15 mm), then 20 ml of MRS which was prepared according to the manufacturers instruction and the procedures described in Cheesbrough (2010), Ekechukwu *et al.* (2025b), Ekesiobi *et al.*, (2025), Ezedianafu *et al.*, (2025a) was added into the plates, allowed to solidified. The plates were incubated in a microaerophilic environment (containing candle used to evacuate all traces of oxygen thereby creating an environment having only carbon iv oxide). The incubation was done for 24 – 48 h at (30±2°C).

Purification of the Isolates

The plate that showed discrete colonies were selected after 24 - 48 h and each colony was aseptically streaked using a sterile wire loop on a sterile poured plate (90mm x 15mm) containing nutrient agar (BIOTECH) prepared according to the manufacturers description. after which it was incubated at their required growth conditions as described by Iheukwumere *et al.* (2020a), Ezedianafu *et al.* (2025b); Idigo *et al.* (2025a), Iheukwumere *et al.* (2025c).

Characterization of the Bacteria Pure Isolates

The pure isolates were characterized using the morphological, biochemical and molecular characteristics as described by Iheukwumere *et al.* (2017a); Iheukwumere *et al.* (2018a), Ike *et al.* (2025a), Iheukwumere *et al.* (2025d).

Morphological characteristics of the Bacteria isolates

The cultural descriptions (size, appearance, edge, elevation, colour) of the isolates were carried out as described in Goldman and Green (2009); Iheukwumere *et al.* (2017b), Iheukwumere *et al.* (2018b), Iheukwumere *et al.* (2020b). The Gram staining technique which revealed the Gram reaction, cell morphology and cell arrangement were also carried out

using the procedure described by Cheesbrough (2010), Goldman and Green (2009) Frank and Robert (2015), Iheukwumere *et al.* (2022b), Iheukwumere *et al.* (2023a). The presence or absence of capsule was also carried out as described by Goldman and Green (2009), Ike *et al.* (2025b), Obiefuna *et al.* (2025a). The presence or absence of flagellum was determined by carrying out motility test as described by Cheesbrough (2010), Iheukwumere *et al.*, (2017c), Iheukwumere *et al.* (2018c), Iheukwumere and Iheukwumere (2022a).

Gram staining technique

A thin smear was made in a cleaned grease free microscopic slide (75mm×25mm), air dried heat fixed. The smear was flooded with crystal violet solution (0.2%) for 60 seconds and rinsed with cleaned water. Gram iodine solution (0.01%) was then applied and allowed for 60 seconds. This was rinsed with cleaned water. This was followed by decolorizing the slide content with 95%w/v ethyl alcohol for 10seconds and then rinsed with cleaned water. The smear was then counter stained with safranin solution (0.025%) for 60 seconds, rinsed with cleaned water, blot drained and air dried. The stained smear was covered with a drop of immersion oil and observed under a binocular compound light microscope using × 100 objective lens as described by Iheukwumere *et al.* (2017d); Iheukwumere *et al.* (2020c), Chude *et al.* (2020), Iheukwumere and Iheukwumere (2022b), Iheukwumere *et al.* (2022c).

Motility test: A semi-solid medium prepared by mixing 5.0g of bacteriological agar (BIOTECH) with 2.0g of nutrient broth (BIOTECH) in 1 Litre of distilled water was used. The solution was dissolved and sterilized using autoclaving technique after dispensing 10 ml portion in different test tubes. The test tubes were allowed to set in vertical positions and then inoculate the test organisms by performing a single stab down the centre of the test tube to half the depth of the medium using sterile stabbing needle. The test tubes were kept in an incubator in vertical position at 35±2°C for 24h (Iheukwumere *et al.*, 2017e; Iheukwumere and Iheukwumere, 2022c; Iheukwumere *et al.*, 2022d; Idigo *et al.*, 2025b).

Biochemical characteristics of the isolates

Indole test: Indole is a nitrogen containing compound formed when the amino acid tryptophan is hydrolyzed by bacteria that have the enzyme tryptophanase. This is detected by using KOVAC's reagent. For this test, isolates were cultured in peptone water in 500.0 ml of deionized water. Ten millilitres of peptone water was dispensed into the test tubes and sterilized. The medium was then inoculated with the isolates and kept in an incubator at 37°C for 48 hr. Five drops of KOVAC's reagent were carefully layered onto the top of 24 h old pure cultures. The presence of indole was revealed by the development of red layer colouration on the top of the broth cultures as described by Iheukwumere *et al.* (2022e), Iheukwumere and Iheukwumere (2022d), Iheukwumere *et al.* (2023b), Egbe *et al.* (2025a), Ike *et al.* (2025c).

Sugar fermentation test: The capability of the isolates to metabolize some sugars (glucose, xylose, ducitol, maltose,

arabinose, inositol, mucate and lactose) with the resulting formation of acid and gas or either were carried out using sugar fermentation test. One litre of 1% (w/v) peptone water was added to 3 mL of 0.2% (w/v) bromocresol purple and 9 ml was dispensed in the test tube that contained inverted Durham tubes. The medium was then sterilized by autoclaving. The sugar solution were prepared at 10% (w/v) and sterilized. One milliliter of the sugar was dispensed aseptically into the test tubes. The medium was then inoculated with the appropriate isolates and the cultures incubated at 37°C for 48 h and were examined for the formation of acid and gas. Change in colour from purple to yellow indicated acid formation while gas formation was assessed by the presence of bubbles in the inverted (Iheukwumere *et al.*, 2022f; Iheukwumere and Iheukwumere, 2022e; Egbe *et al.*, 2025b; Idigo *et al.*, 2025c)

Methyl red test: The glucose phosphate broth was prepared according to the manufacturer's direction and the isolates were aseptically inoculated into the sterilized medium. This was incubated at 37°C for 48 hr. After incubation, five drops of 0.4 % solution of alcoholic methyl red solution was added and mixed thoroughly, and the result was read immediately. Positive tests gave bright red colour while negative tests gave yellow colour (Ezedianafo *et al.*, 2025c; Ike *et al.*, 2025c).

Voges-Proskauer test: The glucose phosphate broth was prepared in accordance to the manufacturer's direction and the isolates were aseptically inoculated into the sterilized medium. This was incubated at 37°C for 48hr. After incubation, 1.0 mL of 40% potassium hydroxide (KOH) containing 0.3% Creatine and 3 ml of 5% solution of α -naphthol was added in the absolute alcohol. Positive reaction was observed by the development of pink colour within five minutes (Egbe *et al.*, 2025b; Ekechukwu *et al.*, 2025c).

Citrate utilization test: The Simmon's Citrate Agar was prepared according to the manufacturer's direction and the isolates were inoculated by stabbing directly at the center of the medium in the test tubes and incubated at 37°C for 48 hr. Positive test was shown by the appearance of growth with blue colour, while negative test showed no growth and the original green colour was retained (Idigo *et al.*, 2025d; Ezedianafo *et al.* 2025d).

Catalase test: The test was carried out as described by Cheesbrough (2010). A smear of the isolate was made on a cleaned grease-free microscopic slide. Then, a drop of 30% hydrogen peroxide (H₂O₂) was added on the smear. Prompt effervescence indicated catalase production (Idigo *et al.*, 2025e; Idigo *et al.*, 2025f).

Oxidase test: The test involved two drops of freshly prepared oxidase reagent dispensed on Whatman No. 1 filter paper which was placed in Petri dish, and a smear of the test isolate was made on the spot using a sterile stick. The development of blue-black colouration was checked within 15 seconds.

Urease test: This was carried out as described by Cheesbrough (2010), Idigo *et al.* (2025g) and Idigo *et al.* (2025h). The urea agar slant was prepared in accordance to the manufacturer's

direction and the isolates were aseptically inoculated into sterilized medium. This was incubated at 37°C for 48 h. After incubation, observation was made for the presence of purple-pink colouration.

Molecular characterization of the isolates

Extraction and purification of DNA: All strains were plated on Nutrient Agar (Biotech) and incubated at 37°C for 24 hr. By means of the procedures of Zymo Research (ZR) DNA miniprep™ kit, bacterial genomic DNA was then extracted and purified (Category No. D6005; Irvine, California, USA) as described by Iheukwumere *et al.* (2018) Iheukwumere *et al.* (2025e; Idigo *et al.*, 2025h).

Determination of the quality of extracted DNA: Using mass spectrophotometer (Nanodrop), One micro litre (1µL) was aseptically dropped into a fresh space in the chamber and the chamber was lightly closed which was then linked to a computer system which showed the window that discovered the value of the sample at 260/280nm as described by (Iheukwumere *et al.*, 2018; Iheukwumere *et al.*, 2025f; Idigo *et al.*, 2025i).

Amplification of DNA and gel electrophoresis of PCR product:This was analysed using Master cycler Nexus Gradient (Eppendorf). A mixture of primer (20 µL), template DNA (20µL), water (72 µL) and master mix (108 µL), which comprises taq polymerase, dimethylsulfoxide (DMSO), magnesium chloride (MgCl₂) and nucleotides triphosphates (NdTPs), was made in 1.5 mL tube and homogenized using vortex mixer (Eppendorf). This was then positioned in the block chamber of the master cycler and then programmed. The PCR program for conditions were as follows: initial incubation at 94°C for 5 mins, followed by 35 cycles of denaturation at 94°C for 15 secs, annealing at 55°C for 15 secs, elongation at 72°C for 21 secs and final extension period for 10 mins at 72°C. The amplified products were electrophoresed in 1.0% agarose gel and a 1kb DNA ladder was used as a size reference. After staining with 3µL of nucleic acid stain (GR green), the gel was documented with gel documentation apparatus (Iheukwumere *et al.*, 2018; Iheukwumere *et al.*, 2025g; Idigo *et al.*, 2025j; Idigo *et al.*, 2025k).

DNA sequencing of 16s rRNA fragment:The 16S rRNA amplified PCR products generated from universal primer (16S), was used for the sequencing using ABI DNA sequencer (Applied Biosystem Inc) at International Institute of Tropical Agriculture (IITA), Ibadan using the method of Iheukwumere *et al.* (2018), Iheukwumere *et al.*, (2025h), and Idigo *et al.* (2025l), Idigo *et al.*, (2025m).

Computational Analysis: This was analysed making use of the modified method of Iheukwumere *et al.* (2018), Iheukwumere *et al.* (2025i), Idigo *et al.* (2025n), Iheukwumere *et al.*, (2025j). The chromatograms generated from the sequences were cleaned to obtain regions with normal sequences. The cleaned nucleotides were aligned using pair wise alignment tool. The consensus sequences formed by the alignment of the forward and reverse sequences were used to perform the Basic Local Alignment Search Tool (BLAST)

using National Centre for Biotechnology Information BLAST over the internet. The sequences of the isolates with 95% and above similarities were accepted. Also the maximum scores, total scores and accession numbers of the isolates were assessed. The relatedness of the isolates was determined by tracing their phylogenetic tree using DNA distance neighbour phylogenetic tree tool.

Preparation of Feed Supplement

Preparation of the corn chaff

The corn chaff was properly collected from the appropriate sites, washed and air dried. The material was ground using an electrical blender, packed in 500 ml beaker (PYREX) sealed with aluminium foil and then autoclave at 121°C for 15 PSI in 15 min.

Fermentation Process

This was carried out using the modified method of Iheukwumere *et al.* (2022), Iheukwumere *et al.* (2025k), Iheukwumere *et al.* (2025l). After autoclaving, a 100 g of the sterile sample was weighed into another 250 ml beaker (PYREX) using analytical weighing balance, which was properly sterilized using electric oven at 180°C for 2 h. This was then inoculated with the fermenter (10 ml) prepared and diluted to a turbidity that matched 0.5 MacFarland standard that was prepared by mixing 0.6mL of 1% BaCl₂. 2H₂O and 99.4 mL of 1% Conc. H₂SO₄. This was allowed for 7 days.

Storage and packaging

After fermentation, the fermented samples were aseptically dried using an electric oven at 80°C for 7days. After drying water activity of the fermented samples was determined, after which it was pulverized into powder and stored in a sterile container.

Moisture Content Determination

A crucible was dried, cooled, and weighed (initial weight recorded as W₁). Then, 2.0 grams of the sample was added to the crucible, and its weight was recorded as W₂. The crucible with the sample was heated in an oven at 105°C for 4 to 6 hours. After heating, the final weight of the crucible and its contents was measured (final weight recorded as W₃). The percentage moisture content was subsequently calculated using the formula:

$$\% \text{ moisture content} = \frac{W_2 - W_3}{W_2 - W_1} \times \frac{100}{1}$$

Experimented Chicks: A total of twenty four (24) broiler chicks (3 weeks old) were purchased from poultry market located at Ihiala market, Ihiala L. G. A. in Anambra State were used for the study. The chicks were kept in separate, thoroughly cleaned and disinfected house and provided with feeds and water ad libitum. All the chicks were vaccinated against Newcastle disease using Lasota vaccine strains at 6 and 19 days of age, against infectious bronchitis using live H120 strain at 6 days old and also against avian influenza (A1) disease using inactivated H5N1 virus vaccine strain at 7 days old. All the vaccines were given via eye drop instillation except (A1) vaccine, which was given through the

subcutaneous route at the back of the neck from the folder report collected from the poultry farmer.

Feed Additive

The fermented groundnut chaff was mixed with fish meal and the feed in a ratio of 1:20. This mixture was properly and thoroughly mixed and administered to the chicks. The chicks were divided into two groups (A and B). Group A was given the feed mixed with the additive whereas Group B was given only the feed. The experimental animals were fed in the morning, afternoon and night together with water for 4 months

Body weights: The body weights of the experimented rats were checked and recorded weekly using electronic weighing balance (LXD200) and recorded as described in the work published by Nwobodo *et al.* (2018), Iheukwumere *et al.* (2025m)

Hematological Indices: The blood samples collected from the broiler chicks were examined using Automated Hematology Analyzer (MIN DRAY BC – 360), and the variations in the red blood cells (RBCs), lymphocytes, monocytes, neutrophils, eosinophils and basophils were assessed and recorded as described in the work published by Agiang *et al.* (2017), Iheukwumere *et al.* (2025n).

Statistical Analysis: The data obtained in this study were presented in tables and figures. Their percentages were also calculated. The sample means and standard deviations of some of the analytical data were also calculated. The significance of this study was determined at 95% using one way analysis of variance (ANOVA). Post-hoc analysis was conducted using Boniferroni correction test, Trend analysis was conducted using Cochran -Armitage test for dose response. Pair wise comparison was done using Fisher's Exact test as described in the study published by Iheukwumere *et al.* (2018), Idigo *et al.*, (2025o), Idigo *et al.* (2025p), Idigo *et al.* (2025q), Idigo *et al.* (2025r), Idigo *et al.* (2025s), Idigo *et al.* (2025t), Manasseh *et al.* (2025).

Results

The cultural and morphological characteristics of the fermenter are shown in Table 1. The result revealed that isolate P showed cream white appearance on MRS agar with a low convex elevation, smooth edge, and surface. The isolate has a Gram-positive rod, non-sporing and non-motile rods. The biochemical characteristics of the fermenters revealed that isolate P was catalase, citrate, oxidase, urase, gelatin, methyl red, Voges Proskauer negative. The isolate showed complex utilization to glucose, lactose, maltose and fructose while it was unable to utilize xylose and sorbitol. The isolate showed varying utilization to D mannitol, inositol, Trehalose and Dulcitol as shown in table 2.

The nucleic acid extracted from the fermenters revealed that it was deoxyribonucleic acid (DNA) at the ratio of 260nm/280nm ranging from 1.80-1.83 as shown in Table 3. The molecular characteristics of the fermenter reveal the presence of *Lactobacillus acidophilus* strain DSM20079

chromosome with a complete genome (LADSM) as shown in Table 4.

The body weight of the chicks when feed with fermented chicken feather revealed that there was a significant ($P < 0.05$) increase in the body weight of the chicks compared to the control group and this was very pronounced at week six of feeding the chicks with the fermented feed supplement as shown in Table 5.

The result of the organ weight of the chicks after been feed with fermented supplement (corn chaff) revealed that there was an increase in the organ weights (Liver, Kidney, Lungs, Heart) of the test chicks compared to the control group and these changes were statistically significant ($P < 0.05$) as shown in Table 6.

The feed conversion ratio and feed intake of the chicks in shown in table 1. The result revealed that during the first week,

that FCR of the test group was higher than the control group, during the second week the FCR of the control group was higher than the test group, and also in week three the FCR of the control group was also higher than the test group same as week four but during week 5 and 6 the FCR of the test group was higher than the control group, also the result revealed that weight and weight gain of the test group was higher than the control group and this indicates that the fermented feed supplement (corn chaff) had a significant effect on the broiler chicks as shown in Table 7.

The blood indices of the chicks fed with the fermented feed supplement (corn chaff) revealed that there was an increase in the counts of WBC, RBC, and in the percentage of monocyte, basophil and lymphocyte of the test group but reduced the number of the PLT, Neutrophil, Eosinophil amongst the test group compared to the control group as shown in Table 8.

Table 1: Cultural and morphological characteristics of the fermenter

Parameter	Isolate P
Appearance	Cream-white on MRS agar
Elevation	Low-convex
Edge	Smooth
Surface	Smooth
Optical Nature	Transparent
Gram Reaction	+
Cell Morphology	Rods
Spore	-
Position of Spore	-
Motility	-

++ Positive; - = Negative

Table 2: Biochemical characteristics of the fermenter

Parameter	Isolate P
Catalase	-
Citrate	-
Oxidase	-
Urease	-
Gelatin	-
Methyl Red	-
Voges Proskauer	-
Glucose	+
D-mannitol	+/_
Lactose	+
Maltose	+
Xylose	-
Inositol	+/_
Fructose	+
Sorbitol	-
Trehalose	+/_
Dulcitol	+/_
Possible Isolate	<i>Lactobacillus</i> species

Table 3: Authentication of nucleic acids extracted from the fermenter

Sample ID	Nucleic Acid Conc($\mu\text{g/mL}$)	260 nm	280 nm	260/280
P	142.40	3.1915	1.7440	1.83

Table 4: Molecular identities of the fermenter

Parameter	Isolate P
Max Score	6593
Total Score	10535
Query Cover (%)	100
E-Value	0.0
Identity (%)	100
Accession Length	2009973
Accession Number	CP020620.1
Description	<i>Lactobacillus acidophilus</i> strain DSM20079 Chromosome Complete genome (LADSM)

Table 5: Body weights of the chicks

Week	Control Group	Test Group
1	168	186
2	376	419
3	660	721
4	974	1052
5	1220	1424
6	1442	1824

Table 6: Organ weight of the chicks

Organ	Control group	Test group
Liver (g)	7.40 ± 0.01	7.41 ± 0.01
Kidney (g)	0.52 ± 0.01	0.51 ± 0.01
Lungs (g)	1.31 ± 0.01	1.30 ± 0.01
Heart (g)	0.67 ± 0.01	0.66 ± 0.01

Table 7: Feed intake and feed conversion ratio among the chicks

Week	Control Group				Test Group			
	Feed (g)	Weight (g)	Weight gain (g)	FCR	Feed (g)	Weight (g)	Weight gain (g)	FCR
1	28	168	60	2.143	34	186	86	2.5294
2	62	376	208	3.355	69	419	223	3.2319
3	102	660	284	2.784	117	721	302	2.5812
4	146	947	287	1.966	158	1052	331	2.0949
5	194	1220	273	1.407	208	1424	372	1.7885
6	243	1442	222	0.914	268	1824	400	1.4925

Table 8: Hematological indices (corn chaff)

Parameter	Control Group	Group fed
WBC (X10 ⁹ L)	12.88	13.68
RBC (X10 ⁹ L)	7.37	7.89
PLT (X10 ⁹ L)	825.00	882.00
Neu (%)	40.60	22.20
Eos (%)	4.75	0.20
Mon (%)	3.85	4.20
Bas (%)	0.10	0.10
Lym (%)	50.70	73.30

Discussion

Fermented feeds have been shown to reduce anti-nutritional factors, improve palatability, increase beneficial microflora in the gut, and enhance the bioavailability of nutrients and when applied to chicken feather, fermentation can improve its amino acid profile, reduce mycotoxin risks, and increase lactic acid production, which benefits gut health and pathogen resistance in poultry (Hur *et al.*, 2014). Similarly, fermented fish meal can further enhance its nutritional value by partially hydrolyzing proteins into peptides and amino acids, making them easier for broilers to digest and absorb (Mohammadi *et al.*, 2021). This present study focuses on the assessment of fermented chicken feather mixed with fish meal as a chicken additive for healthy broiler chicks. The cultural, morphological and biochemical characteristics of the *Lactobacillus* species agree with the findings of many researchers (Tufail *et al.*, 2011; Mannan *et al.*, 2017; Divisekera *et al.*, 2019; and Celik *et al.*, 2021) who isolated and characterized *Lactobacillus* species from yoghurt and banana. The ability of the isolates to utilize common sugars such as glucose, lactose, sucrose, maltose and some alcoholic sugars such as mannitol, sorbitol and xylitol indicates their potential to produce beneficial metabolites which support gut health and aid food fermentation. These potentials were also described by many researchers (Tufail *et al.*, 2011; Mannan *et al.*, 2017; Divisekera *et al.*, 2019; and Celik *et al.*, 2021). The molecular characterization of the *Lactobacillus* species revealed the presence of *Lactobacillus acidophilus* strain DSM20079 (LADSM). The occurrence of *Lactobacillus acidophilus* in strains had been reported by Li *et al.* (2011), Anjum *et al.* (2014), and Mannan *et al.* (2017), who stated that the occurrence may be due to antagonism.

The results of fermented chicken feather showed an increased, weight gain, body weight, organ weight, blood indices and improved FCR and this corroborated the findings of many researchers (Balogu *et al.*, 2017; Onunkwo and Ekine, 2020; and Yang *et al.*, 2020) who evaluated the nutritional quality of fermented corn chaff.

Conclusion

Fermented corn chaff, mediated by *Lactobacillus acidophilus*, significantly enhances growth performance and immune response in broiler chicks, demonstrating its potential as a sustainable feed additive. The study highlights the utilization of agricultural waste, promoting eco-friendly poultry production.

Acknowledgment

We are grateful to all our study participants who join the study voluntarily. We are grateful to ZAHARM Analytical and Research Laboratory, Amawbia, Awka Anambra State, Nigeria for providing enabling environment, resources and techniques for this study. We really salute their wonderful efforts.

Conflict of interests: The authors declare that they have no conflict of interests.

Funding: This research did not receive specific grant from any funding agencies.

References

- Adhikari, P., Kiess, A., Adhikari, R. and Jha, R. (2020). An approach to alternative strategies to control avian coccidiosis and necrotic enteritis. *Journal of Applied Poultry Research* 29:515–534.
- Aljumaah, M.R., Alkhulaifi, M.M., Abudabos, A.M., Alabdullatifb, A., El-Mubarak, A.H. and Al Suliman, A.R. (2020). Organic acid blend supplementation increases butyrate and acetate production in *Salmonella enterica* serovar Typhimurium challenged broilers. *PLoS One* 15: 12 – 17
- Amadi, R.E., Iheukwumere, I.H. and Unaeye, B.C. (2017). Effects Of Crude Alkaloid Extracted From *Ocimum Gratissimum* On The Activity Of Ciprofloxacin Against *Salmonella Enterica* Serovar Typhi. *Advances in Life Science and Technology* 58.
- Anjum, N., Maqsood, S., Masud, T., Ahmad, A., Sohail, A., & Momin, A. (2014). *Lactobacillus acidophilus*: characterization of the species and application in food production. *Critical reviews in food science and nutrition*, 54(9), 1241-1251
- Awad, W.A., Hess, C. and Hess M. (2017). Enteric pathogens and their toxin-induced disruption of the intestinal barrier through alteration of tight junctions in chickens. *Toxins* 9:60 – 66
- Balogu, T. V., Ibrahim, H., & Balogu, D. O. (2017). Nutritionally improved corn mill waste (chaff) with microbial protein: An economic alternative for poultry feed. *Advances in Applied Sciences*, 2(2), 18-22.
- Bell, V., Ferrão, J., & Fernandes, T. (2017). Nutritional guidelines and fermented food frameworks. *Foods*, 6(8), 65. <https://doi.org/10.3390/foods6080065>
- Cabel, M. C., Goodwin, T. L., & Waldroup, P. W. (1988). Feather meal as a nonspecific nitrogen source for abdominal fat reduction in broilers during the finishing period. *Poultry Science*, 67(2), 300-306.
- Cai, Y., Li, Y., Zhang, Y., & Wang, Y. (2019). The role of *Lactobacillus* in immune regulation: Current status and perspectives. *Frontiers in Immunology*, 10, 1804. <https://doi.org/10.3389/fimmu.2019.01804>
- Chachaj, R., Sembratowicz, I., Krauze, M., & Ognik, K. (2019). The effect of partial replacement of soybean meal with fermented soybean meal on chicken performance and immune status. *Journal of Animal and Feed Sciences*, 28(3), 263-271.
- Chen, M., Sun, Q., & Giovannucci, E. (2021). Dairy consumption and risk of type 2 diabetes: The role of fermented dairy products. *American Journal of Clinical Nutrition*, 113(5), 1100–1111. <https://doi.org/10.1093/ajcn/nqaa397>
- Chude, C.O., Iheukwumere, I.H., Iheukwumere, C.M., Nwaolisa, C.N., Egbuna, C., Nwakoby, N.E. and Egbe, P.A. (2020). Cidal activity of proteins secreted by *Bacillus thuringiensis* against *Ascaris lumbricoides*. *International Journal of Research Publications* 49(1): 1033 – 1045.
- Corrêa-Oliveira, R., Fachi, J.L., Vieira, A., Sato, F.T. and Vinolo, M.A.R. (2016). Regulation of immune cell function by short-chain fatty acids. *Clinical Translational Immunology* 5:73 – 80
- Cristofori, F., Dargenio, V. N., Dargenio, C., Miniello, V. L., Barone, M., & Francavilla, R. (2021). Anti-inflammatory and immunomodulatory effects of probiotics in gut inflammation: A door to the future. *Frontiers in Immunology*, 12, 578386. <https://doi.org/10.3389/fimmu.2021.578386>
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025a). Multiple Antibiotic Resistance Bacterial Strains in Frozen Meat Sold at Abagana, Anambra State: A Public Health Concern. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 181–186. <https://doi.org/10.54117/ijamb.v4i3.75>
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025b). The Burden of Antibiotic Resistance: Evaluating the Impact of Multiple Antibiotic-Resistant Enteric Bacteria in Academic Environments. *IPS Interdisciplinary Journal of Biological Sciences*, 4(4), 144–149. <https://doi.org/10.54117/ijbbs.v4i4.78>
- Dim, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Ugwu, C. H., Ike, V. E., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025c). Antimicrobial resistance in aquaculture: evaluating *Pseudomonas aeruginosa* from fish ponds. *IPS Intelligentia Multidisciplinary Journal*, 4(1), 32–36. <https://doi.org/10.54117/iimj.v4i1.10>

- Dimidi, E., Cox, S. R., Rossi, M., & Whelan, K. (2019). Fermented foods: Definitions and characteristics, impact on the gut microbiota and effects on gastrointestinal health and disease. *Nutrients*, 11(8), 1806. <https://doi.org/10.3390/nu11081806>
- Dittoe, D.K., Ricke, S.C., Kiess, A.S. (2018). Organic acids and potential for modifying the avian gastrointestinal tract and reducing pathogens and disease. *Frontiers in Veterinary Science* 5:216 – 223
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., & Ezeumeh, E. N. (2025a). Antibiotic Susceptibility of *Helicobacter pylori* Isolates from Patients at Nnewi Teaching Hospital, Anambra State. *IPS Journal of Basic and Clinical Medicine*, 2(2), 51–57. <https://doi.org/10.54117/ijbcm.v2i2.11>
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., Ezeumeh, E. N., & Egbuna, C. (2025b). *Helicobacter pylori* Inhibition by Medicinal Plant Extracts: An In Vitro Assessment. *IPS Journal of Drug Discovery Research and Reviews*, 3(1), 32–37. <https://doi.org/10.54117/ijddr.v3i1.28>
- Egbe, P. A., Umeaku, C. N., Iheukwumere, I. H., Iheukwumere, C. M., Onwuasoanya, U. F., Ezenwata, I. S., Afulukwe, S. C., Ike, V. E., & Ezeumeh, E. N. (2025c). Medicinal Plant Extracts Enhance Conventional Antibiotic Activity against *Helicobacter pylori*: An In Vitro Assessment. *IPS Interdisciplinary Journal of Biological Sciences*, 4(2), 93–99. <https://doi.org/10.54117/ijbs.v4i2.51>
- Ejike, C.E., Iheukwumere, I.H. and Armadi, R.E. (2017). Susceptibility of *Escherichia coli* Isolated from Oligospermia Patient to *Gongronema latifolium* leaves extract. *J. Biol. Agriculture. Healthcare* 7(14).
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025a). Bacterial Loads of Smoked Fish and Chicken: Role of pH and Moisture Content. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 3(1), 44–49. <https://doi.org/10.54117/iajnfms.v3i1.102>
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025b). Biological Inhibition of Pathogenic Bacteria Isolated from Smoked Fish and Chicken: An In Vitro Study. *IPS Interdisciplinary Journal of Biological Sciences*, 4(2), 85–92. <https://doi.org/10.54117/ijbs.v4i2.50>
- Ekechukwu, C. C., Umeh, S. O., Iheukwumere, I. H., & Iheukwumere, C. M. (2025c). Prophylactic Potential of the Most Potent Synergistic Biological Agent against Bacterial Infections from Smoked Fish and Chicken. *IPS Journal of Applied Microbiology and Biotechnology*, 4(2), 153–160. <https://doi.org/10.54117/ijamb.v4i2.57>
- Ekesiobi, A. O., Iheukwumere, C. M., Iheukwumere, I. H., Ejike, C. E., Ihechukwu, C. C., Ike, V. E., Okereke, F. O., & Ochibulu, S. C. (2025). Hyping the Inhibitory Activity of *Xylopiya aethiopica* against *Vibrio cholerae* using Azithromycin. *IPS Journal of Basic and Clinical Medicine*, 2(3), 93–98. <https://doi.org/10.54117/ijbcm.v2i3.16>
- Ezedianafo, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025a). *Musca domestica*: A vector of multidrug-resistant enteric bacteria. *Journal of Veterinary, Allied, and One Health Sciences*, 1(2), 30–38. <https://doi.org/10.54117/3vwy0p36>
- Ezedianafo, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025b). Occurrences of meropenem- and imipenem-resistant *Klebsiella pneumoniae* in *Musca domestica* in hospital landfills. *African Journal of Applied Research & Sustainable Development*, 1(2), 25–35. <https://doi.org/10.54117/wjmnvy91>
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025). Multiple antibiotic resistance indices of enteric bacteria isolated from *Musca domestica*. *Journal of Public Health, Policy, and Society*, 1(2), 29–37. <https://doi.org/10.54117/k8r78723>
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025). *Klebsiella pneumoniae* isolated from *Musca domestica*: Antibiotic susceptibility and resistance patterns. *Journal of Veterinary, Allied, and One Health Sciences*, 1(2), 39–47. <https://doi.org/10.54117/vy6y8f94>
- Ezendianefor, J. N., Iheukwumere, I. H., Iheukwumere, C. M., Okolo, O., Nwike, I., & Ubajekwe, C. C. (2025). *Musca domestica* as vectors of pathogenic enteric bacteria: A public health concern. *African Journal of Applied Research & Sustainable Development*, 1(2), 36–45. <https://doi.org/10.54117/s671mk28>
- Frias, J., Peñas, E., & Martínez-Villaluenga, C. (2017). Fermented foods in health promotion and disease prevention. In V. R. Preedy & R. R. Watson (Eds.), *Fermented Foods in Health and Disease Prevention* (pp. 3–20). Academic Press.
- Gharib-Naseri, K., Dorigam, J.C., Doranalli, K., Morgan, N., Swick, R.A. and Choct M. (2021). *Bacillus amyloliquefaciens* CECT 5940 improves performance and gut function in broilers fed different levels of protein and/or under necrotic enteritis challenge. *Animal Nutrition* 7:185–197
- Gharib-Naseri, K., Kheravii, S., Keerqin, C., Morgan, N., Swick, R. and Choct, M. (2019). Two different *Clostridium perfringens* strains produce different levels of necrotic enteritis in broiler chickens. *Poultry Science* 98:6422–6432
- Guo, Q., Goldenberg, J. Z., Humphrey, C., El Dib, R., Johnston, B. C., & Lo, C. K. F. (2020). Probiotics for the prevention of pediatric antibiotic-associated diarrhea. *Cochrane Database of Systematic Reviews*, 4(4), CD004827. <https://doi.org/10.1002/14651858.CD004827.pub5>
- Guo, S., Zhang, Y., Cheng, Q., Xu, J., Hou, Y., Wu, X., Du, E., & Ding, B. (2020). Partial Substitution of Fermented Soybean Meal for Soybean Meal Influences the Carcass Traits and Meat Quality of Broiler Chickens. *Animals*, 10(2), 225. <https://doi.org/10.3390/ani10020225>
- Hill, C., Guarner, F., Reid, G., Gibson, G. R., Merenstein, D. J., Pot, B., ... and Sanders, M. E. (2014). Expert consensus document: The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nature Reviews Gastroenterology and Hepatology*, 11(8), 506–514. <https://doi.org/10.1038/nrgastro.2014.66>
- Husnain, A., Anwar, U., Fatima, A., Mustafa, R., Farooq, U., Abbas, W., Khalid, M. F., Ashraf, M., & Aziz ur Rahman, M. (2025). Effects of replacement of soybean meal with fermented soybean meal on growth performance, nutrient digestibility and carcass characteristics in broiler. *Livestock Science*, 299, 105779. <https://doi.org/10.1016/j.livsci.2025.105779>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Bacterial symbionts of *Rhynchophorus phoenicis*: Modulation of rat lymphocyte function and immunity. *Journal of Tropical Medicine and Public Health Solutions*, 1(2), 55–63. <https://doi.org/10.54117/jtmphs.v1i2.36>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Multidrug-resistant *Klebsiella pneumoniae* in *Musca domestica*: A potential public health threat. *Journal of Tropical Medicine and Public Health Solutions*, 1(2), 46–54. <https://doi.org/10.54117/jtmphs.v1i2.35>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Termite-derived antimicrobials: A novel approach to control *Bacillus cereus* in food products. *International Journal of Global Trends and Research*, 1(2), 45–57. <https://doi.org/10.54117/2mgwn270>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Bacterial diversity in insect guts and their potential applications in industry. *International Journal of Global Trends and Research*, 1(2), 36–44. <https://doi.org/10.54117/2k0d9t18>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Bioactive compounds from *Macrotermes*: A novel approach to combat multidrug resistant *Klebsiella pneumoniae* from *Musca domestica* in hospital landfills. *IPS Journal of Biotechnology and Applied Biochemistry*, 1(2), 82–92. <https://doi.org/10.54117/ijbab.v1i2.111>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Assessment of blood indices in rats supplemented with *Macrotermes*-derived nutrient-enhanced bacteria. *IPS Journal of Basic and Clinical Medicine*, 2(4), 143–151. <https://doi.org/10.54117/ijbcm.v2i4.39>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Exploring the potential of termite gut bacteria as growth promoters and organ function enhancers in albino Wistar rats. *Health Science Research International*, 1(2), 43–52. <https://doi.org/10.54117/hsri.v1i2.38>

- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Assessment of hematological parameters in albino Wistar rats fed with *Rhynchophorus phoenicis* larvae enriched with autochthonous bacteria. *Health Science Research International*, 1(2), 34–42. <https://doi.org/10.54117/hsri.v1i2.37>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Entomopathogenic bacteria-mediated management of *Delia radicum*: A step towards eco-friendly pest control. *Journal of Agriculture, Food Technology and Sustainability*, 2(2), 112–121. <https://doi.org/10.54117/jafts.v2i2.110>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Antimicrobial activity of *Macrotermes*-derived eluates against multidrug resistant *Pseudomonas* species: Implications for aquaculture disease management. *Journal of Agriculture, Food Technology and Sustainability*, 2(2), 102–111. <https://doi.org/10.54117/jafts.v2i2.109>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Exploring the potential of entomopathogenic bacteria for sustainable management of *Sitophilus zeamais* in maize storage systems. *IPS Journal of Plant, Animal, and Environmental Sciences*, 1(1), 11–20. <https://doi.org/10.54117/ijpae.v1i1.108>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Biological control of *Acanthoscelides obtusus* using entomopathogenic bacteria. *IPS Journal of Plant, Animal, and Environmental Sciences*, 1(1), 1–10. <https://doi.org/10.54117/ijpae.v1i1.107>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Modulation of phagocytic index in albino Wistar rats via bacterial symbionts from *Rhynchophorus phoenicis* larvae. *African Journal of Nutrition and Applied Research*, 1(2), 27–36. <https://doi.org/10.54117/fm3vgt16>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Bacterial symbionts of insect larvae: A novel approach to improving micronutrient content. *African Journal of Nutrition and Applied Research*, 1(2), 27–36. <https://doi.org/10.54117/960k2266>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Unlocking the potential of termite gut microbiome: Enhancing nutritional value through bacterial symbionts. *IPS Journal of Nutrition and Food Science*, 5(1), 636–645. <https://doi.org/10.54117/ae6gj081>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Enhancing nutritional parameters with bacterial symbionts from *Macrotermes* species: A potential frontier in nutritional biotechnology. *IPS Journal of Nutrition and Food Science*, 5(1), 625–635. <https://doi.org/10.54117/a2b7jb52>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Bacterial symbionts of *Macrotermes* species: Assessing their impact on phagocytic indices of albino Wistar rats. *IPS Interdisciplinary Journal of Biological Sciences*, 5(1), 187–196. <https://doi.org/10.54117/ijbbs.v5i1.106>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Bacterial symbionts of insects: Exploring their role in insect nutritional composition. *IPS Interdisciplinary Journal of Biological Sciences*, 5(1), 177–186. <https://doi.org/10.54117/ijbbs.v5i1.105>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Augmenting rat lymphocyte function by bacterial symbiont of *Macrotermes* species. *IPS Journal of Applied Microbiology and Biotechnology*, 5(1), 281–290. <https://doi.org/10.54117/ijamb.v5i1.104>
- Idigo, M. A., Iheukwumere, I. H., Iheukwumere, C. M., Nnaeze, B. C., Akulue, C. J., Nwakoby, N. E., Ezendianefor, J. N., Ike, V. E., Nnaeozie, A. O., Ezekwueche, S. N., Anagor, I. S., Aniekwe, C. C., Ezeoke, F. C., Okereke, F. O., & Ochibulu, S. C. (2025). Antimicrobial peptides from insects: A study on their efficacy against pathogens. *IPS Journal of Applied Microbiology and Biotechnology*, 5(1), 271–280. <https://doi.org/10.54117/ijamb.v5i1.103>
- Iheukwumere, C. M., & Iheukwumere, I. H. (2022a). Nutritive and Antinutrient Values of Soybean Condiments Produced from Indigenous Fermenters. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 1(1): 1-5. <https://doi.org/10.54117/iajnfms.v1i1.8>
- Iheukwumere, C. M., & Iheukwumere, I. H. (2022e). Hematological indices and sensory quality of fermented soybean condiments. *World Journal of Advanced Research and Reviews*, 14(2), 435-42
- Iheukwumere, C. M., Ekesiobi, A. O., Iheukwumere, I. H., Ejike, C. E., Ilechukwu, C. C., Dim, C. N., & Ochibulu, S. C. (2025g). Dual Approach Therapy: Assessing *Xylopiia aethiopia* and Ciprofloxacin Synergy against *Salmonella enterica* Serovar Typhi. *IPS Intelligentia Multidisciplinary Journal*, 4(1), 27–31. <https://doi.org/10.54117/iiimj.v4i1.9>
- Iheukwumere, C. M., Ekesiobi, A. O., Iheukwumere, I. H., Ejike, C. E., Ilechukwu, C. C., Dim, C. N., Ochibulu, S. C., Unegbu, C. C., & Egbuna, C. (2025h). Food Safety Implications: Assessing the Potential of *Desmodium velutinum* Leaves Extracts to Control the Most Predominant Fungal Contamination in Ready-To-Eat Fried Chicken. *IPS Journal of Nutrition and Food Science*, 4(3), 494–500. <https://doi.org/10.54117/ijnfs.v4i3.111>
- Iheukwumere, C. M., Iheukwumere, I. H., Nwakoby, N. E., Idigo, M. A., & Ike, V. E. (2025m). Evaluation of fermented chicken feather meal as a dietary supplement on rat lipid metabolism and immune response. *African Journal of Nutrition and Applied Research*, 1(1), 17–26. <https://doi.org/10.54117/qbwbbp15>
- Iheukwumere, C. M., Iheukwumere, I. H., Okoli, U. O., & Ugwu, C. H. (2023a). Immunological Impact of Fermented Soybean Condiments Produced from Indigenous Fermenters. *Journal of Advances in Microbiology* 23(10): 27-37
- Iheukwumere, C. M., Iheukwumere, I. H., Ugwu, C. H., & Okoli, U. O. (2023b). Toxicity of Prepared Fermented Soybean Condiments from Indigenous Fermenters. *Journal of Advances in Microbiology* 23(10): 38 – 51.
- Iheukwumere, C. M., Umeaku, C. N., Chukwura, E. N., & Iheukwumere, I. H. (2022f). Characterization of the indigenous fermenters for the production of fermented condiments from soybean seeds. *World Journal of Advanced Research and Reviews*, 14(2), 423-434.
- Iheukwumere, I. H., & Ejike, C. E. (2017b). Comparative study of the inhibitory activities of *Ocimum gratissimum* and *Nepeta cataria* against *Salmonella enterica* serovar Typhi and their larvicidal effect against *Anopheles gambiae*. *African Journal of Education, Science and Technology (AJEST)*, 3(4), 16-24
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnaeozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025c). Exploring the Phytochemical and Antimicrobial Properties of Fruit Vinegar: A Study on *Phoenix Dactylifera* and *Malus Sylvestris*. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 115–122. <https://doi.org/10.54117/ijamb.v4i1.48>
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnaeozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025d). Microbial Quality and Sensory Assessment of Vinegar from Date Palm and Apple Fruits: Implications for Consumer Preference. *IPS Journal of Nutrition and Food Science*, 4(2), 410–417. <https://doi.org/10.54117/ijnfs.v4i2.100>
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnaeozie, C. H., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025f). Safety Evaluation of Vinegar from *Phoenix Dactylifera* and *Malus*

- Sylvestris: Toxicity and Acetic Acid Content. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 123–131. <https://doi.org/10.54117/ijamb.v4i1.49>
- Iheukwumere, I. H., Ajeh, J. C., Iheukwumere, C. M., Ike, V. E., Obianom, A. O., Ihenatuoha, U. A., Igboanugo, E. U., Onwuasoanya, U. F., Okereke, F. O., Nnadozie, C. H., Agbaugo, C. F., Nwike, M. I., Nwakoby, N. E., & Ilechukwu, C. C. (2025k). Exploring the Phytochemical and Antimicrobial Properties of Fruit Vinegar: A Study on Phoenix Dactylifera and Malus Sylvestris. *IPS Journal of Applied Microbiology and Biotechnology*, 4(1), 115–122. <https://doi.org/10.54117/ijamb.v4i1.48>
- Iheukwumere, I. H., Amadi, E. R., & Chude, C. (2018b). Synergistic Effects of Probiotics and Autogenous Bacterin Against Inositol Negative Motile Salmonella Species. *Journal of Biology, Agriculture and Healthcare* 8(6).
- Iheukwumere, I. H., Amadi, R. E., Unaeze, B. C., & Campus, N. (2017c). Enterotoxigenicity Profile of Salmonella Enterica Serovar Typhimurium in Suckling Albino Mice. *Journal of Natural Sciences Research* 7(14).
- Iheukwumere, I. H., Chukwura, E. I., & Chude, C. (2018c). In vivo activities of some selected antimicrobial agents against enteric bacteria isolated from chicken feeds on broiler layers. *Journal of Biology, Agriculture and Healthcare*, 8(6).
- Iheukwumere, I. H., Ejike, C. E., & Okeke, C. E. (2017d). A trial to prevent sorbitol negative Escherichia coli infections in chicks using autogenous bacteria and probiotics. *Journal of Natural Sciences Research*, 7, 56–63.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Ihenatuoha, U. A. (2025a). Cross-Sectional Study of Different Strains of Bacillus cereus among Pap Sold in Major Towns in Ihiala LGA, Anambra State. *IPS Journal of Public Health*, 5(2), 199–204. <https://doi.org/10.54117/ijph.v5i2.39>.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Destiny, E. C. (2025b). Cross-Sectional Study of Major Strains of Salmonella enterica Subspecies Enterica Serovar Typhi among Borehole Used in Uli Community. *IPS Journal of Public Health*, 5(2), 205–210. <https://doi.org/10.54117/ijph.v5i2.40>.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., Udeagbara, O. E., Unaeze, B. C., Obiefuna, O. H., Ike, V. E., Onyemekara, N. N., & Ihenatuoha, U. A. (2025e). Quotidian of Substantial Strain of Shigella dysenteriae among Ready To-Eat Fruit Salad Sold in Uli Community. *Journal of Pollution Monitoring, Evaluation Studies and Control*, 4(1), 95–99. <https://doi.org/10.54117/jpmesc.v4i1.17>
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Ihenatuoha, U. A. (2025i). Cross-Sectional Study of Different Strains of Bacillus cereus among Pap Sold in Major Towns in Ihiala LGA, Anambra State. *IPS Journal of Public Health*, 5(2), 199–204. <https://doi.org/10.54117/ijph.v5i2.39>.
- Iheukwumere, I. H., Iheukwumere, C. M., Obianom, A. O., Nnadozie, C. H., Okereke, F. O., Onwuasoanya, U. F., ... Destiny, E. C. (2025j). Cross-Sectional Study of Major Strains of Salmonella enterica Subspecies Enterica Serovar Typhi among Borehole Used in Uli Community. *IPS Journal of Public Health*, 5(2), 205–210. <https://doi.org/10.54117/ijph.v5i2.40>.
- Iheukwumere, I. H., Iheukwumere, M. C., & Nwakoby, N. E. (2022d). Sequential Pathogenicity Study of SOR+ and SOR-Escherichia coli Isolated from Roasted Meat. *IPS Intelligentsia Multidisciplinary Journal*, 1(1), 1–11.
- Iheukwumere, I. H., Nwankwo, A. K., Iheukwumere, C. M., Okorie, N. A., Nwakoby, N. E., Ekesiobi, A. O. and Okolo, O. C. (2025n). Characterization and pathogenic profile of *Aspergillus fumigatus* isolated from lendfills. *Microbes and Infectious Diseases* 6(4)
- Iheukwumere, I. H., Obi, P. C. and Unaeze, B. C. (2017a). A trial to prevent *Vibrio cholerae* in albino mice using autogenous bacterin. *Advances in Life Science and Technology* 58:34–42
- Iheukwumere, I. H., Uneze, B. C., & Ejike, C. E. (2017e). Efficacy of some selected antimicrobial substances in prevention of enteric bacterial infection in broiler chicks. *J. Biol. Agriculture. Healthcare*, 7, 58–66.
- Iheukwumere, I.H. , Iheukwumere, C.M. , Nnadozie, H. C. ,Unaeze, C.B. , Obiefuna, O.H. Obianom, A.O. and Ejike, C. E. (2024). Hematotoxicological and mosquito larvicidal studies of crystal proteins secreted by Bacillus thuringiensis and Bacillus sphaericus. *Tropical Journal of Applied Natural Sciences* 2(2): 61 – 92.
- Iheukwumere, I.H. and Iheukwumere, M.C. (2022c). Streptococcus suis in Pigs and Environs: A Cross-sectional Study. *IPS Journal of Public Health*, 1(2), 9-12. <https://doi.org/10.54117/ijph.v1i2.4>.
- Iheukwumere, I.H. and Iheukwumere, M.C. (2022g). Cross-sectional Study of Multiple Antibiotic-resistant Streptococcus suis in Pigs and Environments. *IPS Interdisciplinary Journal of Biological Sciences*, 1(1), 19–21. <https://doi.org/10.54117/ijbbs.v1i1.4>
- Iheukwumere, I.H., Dimejesi, S.A., Iheukwumere, C.M., Chude, C.O., Nwaolisa, C.N., Ukoha, C.C., Nwakoby, N.E., Egbuna, C. and Egbe, P.A. (2020) Diversity and molecular characterization of keratinophilic fungi from soil samples. *International Journal of Research Publication* 50(1); 1047 -1062.
- Iheukwumere, I.H., Iheukwumere, M.C. and Nwakoby, N.E. (2022b). Synergistic Effects of Probiotics and Autogenous Bacterin against Salmonella enterica Serovar Typhimurium Strain U288. *IPS Journal of Nutrition and Food Science*, 1(1), 1–5. <https://doi.org/10.54117/ijnfs.v1i1.3>.
- Iheukwumere, I.H., Nwike, M. I., Iheukwumere, C.M., Ike, V.E., Obianom, A.O., Ihenatuoha, U.A., Igboanugo, E.U., Ekesiobi, A.O., Okereke, F.O., Obiefuna, O. H. Nnadozie, C.H., Agbaugo, C.F., Oduoye, O.T., Nwakoby, N.E., Ilechukwu, C. C., Ochibulu, S. C. and Ejike, C. E. (2025l). Extraction and Elucidation of Antibiotics from the Mycelia of *Aspergillus niger* Isolated from Poultry Farm against Enteric Bacterial Pathogens. *IPS Journal of Advanced and Applied Biochemistry*, 1(1), 1–10. <https://doi.org/10.54117/ijaab.v1i1.58>.
- Iheukwumere, I.H., Iheukwumere, C.M., Chude, C.O., Nwaolisa, C.N. and Egbe, P.A. (2020a). Comparative study of different clinical samples used for the diagnosis of staphylococcal systemic infections in apparent healthy students. *International Journal of Research Publications* 49(1): 1 – 10
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025a). Prevalence of Bacillus cereus in Powdered Soybean Sold in Uli Community, Anambra State: A Cross-Sectional Study. *IPS Journal of Basic and Clinical Medicine*, 2(3), 108–114. <https://doi.org/10.54117/ijbcm.v2i3.18>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025b). *Bacillus cereus* in Uli's cornflour: A prevalence study. *IPS Journal of Nutrition and Food Science*, 4(3), 544–548. <https://doi.org/10.54117/8bte840>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025c). Pathogenic Profile Analysis: In Vitro Screening of Enteric Bacteria from University Dusters. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 187–191. <https://doi.org/10.54117/ijamb.v4i3.76>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025d). Frozen Fish Pathogens: Antimicrobial Resistance and Public Health Implications. *IPS Interdisciplinary Journal of Biological Sciences*, 4(4), 138–143. <https://doi.org/10.54117/ijbbs.v4i4.77>
- Ike, V. E., Iheukwumere, I. H., Iheukwumere, C. M., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., & Ochibulu, S. C. (2025e). Stream water quality assessment: Antibiotic resistance of Lac-positive enteric bacterial isolates. *Journal of Pollution Monitoring, Evaluation Studies and Control*, 4(2), 120–125. <https://doi.org/10.54117/jpmesc.v4i2.21.2025>
- Ke, K., Sun, Y., He, T., Liu, W., Wen, Y., Liu, S., ... & Gao, X. (2024). Effects of feather hydrolysates generated by probiotic *Bacillus licheniformis* WHU on gut microbiota of broiler and common carp. *Journal of Microbiology*, 62(6), 473–487.
- Khan, S.H. and Iqbal, J. (2016). Recent advances in the role of organic acids in poultry nutrition. *Journal of Applied Animal Research* 44:359–369
- Kim, J., Lee, H., & Kim, Y. (2018). Effects of fermented milk products on cholesterol and blood pressure: A review of clinical trials. *Journal of Dairy Science*, 101(4), 2724–2741. <https://doi.org/10.3168/jds.2017-13708>
- Kim, S. K., Kim, T. H., Lee, S. K., Chang, K. H., Cho, S. J., Lee, K. W., & An, B. K. (2016). The Use of Fermented Soybean Meals during Early Phase Affects Subsequent Growth and Physiological Response in Broiler Chicks. *Asian-Australasian Journal of Animal Sciences*, 29(9), 1287. <https://doi.org/10.5713/ajas.15.0653>
- Latorre, J.D., Adhikari, B., Park, S.H., Teague, K.D., Graham, L.E. and Mahaffey, B.D.. (2018). Evaluation of the epithelial barrier function and ileal microbiome in an established necrotic enteritis challenge model in broiler chickens. *Frontier in Veterinary Science* 5:199 – 206
- Le Doare, K., Holder, B., Bassett, A., & Pannaraj, P. S. (2020). Mother's milk: A purposeful contribution to the development of the infant microbiota and immunity. *Frontiers in Immunology*, 11, 362. <https://doi.org/10.3389/fimmu.2020.00362>

- Li, Y., Guo, B., Wu, Z., Wang, W., Li, C., Liu, G., & Cai, H. (2020). Effects of Fermented Soybean Meal Supplementation on the Growth Performance and Cecal Microbiota Community of Broiler Chickens. *Animals*, 10(6), 1098. <https://doi.org/10.3390/ani10061098>
- Lv, W., Ma, Y., Zhang, Y., Wang, T., Huang, J., He, S., Du, H., & Guo, S. (2023). Effects of *Lactobacillus plantarum* fermented Shenling Baizhu San on gut microbiota, antioxidant capacity, and intestinal barrier function of yellow-plummed broilers. *Frontiers in Veterinary Science*, 10, 1103023. <https://doi.org/10.3389/fvets.2023.1103023>
- Manasseh, C.O., Logan, C.S.P., Ikeyi, A.P., Ede, K.K., Iheukwumere, I.H., Iheukwumere, C.M. and Ejike, C.E. (2025). Investigating the Effects of the Covid-19 Pandemic and Climate Risks on Trade Balance in Emerging Markets. *The Nigerian Health Journal* 25(2): 1-27. <https://doi.org/10.71637/tnhj.v25i2.914>
- Marco, M. L., Sanders, M. E., Gänzle, M., Arrieta, M. C., Cotter, P. D., De Vuyst, L., ... & Hutkins, R. (2021). The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on fermented foods. *Nature Reviews Gastroenterology and Hepatology*, 18(3), 196–208. <https://doi.org/10.1038/s41575-020-00390-5>
- McKnight, L., Page, G. and Han, Y. (2020). Effect of replacing in-feed antibiotics with synergistic organic acids, with or without trace minerals and/or water acidification, on growth performance and health of broiler chickens under a *Clostridium perfringens* type A challenge. *Avian Disease* 64(3):374–378
- Melini, F., Melini, V., Luziatelli, F., Ficca, A. G., & Ruzzi, M. (2019). Health-promoting components in fermented foods: An up-to-date systematic review. *Nutrients*, 11(5), 1189. <https://doi.org/10.3390/nu11051189>
- Meyer, B., Bessei, W., Vahjen, W., Zentek, J., & Harlander-Matauschek, A. (2012). Dietary inclusion of feathers affects intestinal microbiota and microbial metabolites in growing Leghorn-type chickens. *Poultry Science*, 91(7), 1506–1513. <https://doi.org/10.3382/ps.2011-01786>
- Nwike, M.I., Iheukwumere, I.H. and Uneze, B.C. (2017). Effect of Spices, pH and Temperature on the Survival and Multiplication of *Staphylococcus aureus* in Locally Made Soya Milk Drink. *Journal of Natural Sciences Research* 7(4).
- Obianom, A.O., Iheukwumere, I.H., Iheukwumere, C.M., Ochibulu, S.C., Nnadozie, H. C. and Ifenetu, F. C. (2024). Supersizing the inhibitory activity of *Xylopiya aethiopia* extract against *Vibrio cholerae* using doxycycline. *Tropical Journal of Applied Natural Sciences* 2(2).
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025a). Assessing the Impact of Glycemic Index on Microbial Quality and Storage Stability of Tomato Jam. *IPS Journal of Applied Microbiology and Biotechnology*, 4(3), 192–202. <https://doi.org/10.54117/ijamb.v4i3.81>
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025b). Physicochemical and nutritional properties of tomato jam: Influence of sweetener type and glycemic index. *IPS Journal of Nutrition and Food Science*, 4(3), 561–568. <https://doi.org/10.54117/yazv0114>
- Obiefuna, U. G., Umeh, S. O., & Iheukwumere, I. H. (2025c). Micronutrient profile and acceptability of tomato jam: A comparison of high and low glycemic carbohydrate index sweeteners. *IPS Applied Journal of Nutrition, Food and Metabolism Science*, 3(2), 67–74. <https://doi.org/10.54117/pr4r6c73>
- Onrust, L., Van, Driessche, K., Ducatelle, R., Schwarzer, K., Haesebrouck, F. and Van Immerseel, F. (2018). Valeric acid glyceride esters in feed promote broiler performance and reduce the incidence of necrotic enteritis. *Poultry Science* 97:2303–2311.
- Onunkwo, D. N., & Ekine, O. A. (2020). Performance of broiler chickens fed diet containing fermented maize milling waste. *Nigerian Journal of Animal Production*, 47(1), 214–220.
- Park, K. Y., Jeong, J. K., Lee, Y. E., & Daily, J. W. (2021). Health benefits of kimchi (Korean fermented vegetables) as a probiotic food. *Journal of Medicinal Food*, 27(3), 365–375. <https://doi.org/10.1089/jmf.2020.0155>
- Premathilaka, K. T., Nawarathne, S. R., Nambapana, M. N., Macelline, S. P., Wickramasuriya, S. S., Ang, L., Jayasena, D. D., & Heo, J. M. (2020). Partial or complete replacement of fishmeal with fermented soybean meal on growth performance, fecal composition, and meat quality in broilers. *Journal of Animal Science and Technology*, 62(6), 824. <https://doi.org/10.5187/jast.2020.62.6.824>
- Safari, H., & Mohit, A. (2024). Feather meal processing methods impact the production parameters, blood biochemical indices, gut function, and hepatic enzyme activity in broilers. *Journal of Animal Science*, 102. <https://doi.org/10.1093/jas/skae068>
- Safari, H., Mohit, A., & Mohiti-Asli, M. (2024). Fermented feather meal improves the antioxidant status, meat quality, and immune response of broilers. *Iran J Vet Med*, 1-43.
- Salehizadeh, M., Ebrahimi, M. T., Mousavi, S. N., Sepahi, A. A., & Orooji, R. (2025). Transforming Feather Meal Into a High-Performance Feed for Broilers. *Veterinary Medicine and Science*, 11(1), e70199. <https://doi.org/10.1002/vms3.70199>
- Savaiano, D. A., & Hutkins, R. W. (2020). Yogurt, cultured fermented milk, and health: A review. *Nutrients*, 12(5), 1256. <https://doi.org/10.3390/nu12051256>
- Soumei, E., Mohebodini, H., Toghyani, M., Shabani, A., Ashayerizadeh, A., & Jazi, V. (2019). Synergistic effects of fermented soybean meal and mannan-oligosaccharide on growth performance, digestive functions, and hepatic gene expression in broiler chickens. *Poultry Science*, 98(12), 6797–6807. <https://doi.org/10.3382/ps/pez409>
- Stanton, C., Ross, R. P., Fitzgerald, G. F., & van Sinderen, D. (2021). Fermented functional foods: Trends and challenges. *Current Opinion in Food Science*, 40, 114–122. <https://doi.org/10.1016/j.cofs.2021.02.005>
- Sun, Y., Han, Y., Chen, J., Ni, A., Jiang, Y. and Li Y. (2020). Effect of replacing in-feed antibiotics with synergistic organic acids on growth performance, health, carcass, and immune and oxidative status of broiler chickens under *Clostridium perfringens* type A challenge. *Avian Disease* 64:393–400
- Ugwu, C. H., Iheukwumere, I. H., Iheukwumere, C. M., Ike, V. E., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025a). Maternal health and antibiotic resistance: *Klebsiella pneumoniae* isolates analysis. *IPS Journal of Public Health*, 5(3), 290–295. <https://doi.org/10.54117/s3tx6v26>
- Ugwu, C. H., Iheukwumere, I. H., Iheukwumere, C. M., Ike, V. E., Dim, C. N., Ezendianefo, J. N., Egbe, P. A., Oragwu, I. P., Orji, C. C., Ogbonnaya, O. C., Onwuasoanya, U. F., Okereke, F. O., Oduenyi, P. M., & Ochibulu, S. C. (2025b). *Ocimum gratissimum* Extract's Effectiveness against *Vibrio cholerae* from Uli Streams. *IPS Journal of Phytochemistry and Medicinal Plant Research*, 1(2), 15–19. <https://doi.org/10.54117/ijpmpr.v1i2.38>
- Wang, J., Yao, L., Su, J., Fan, R., Zheng, J., & Han, Y. (2023). Effects of *Lactobacillus plantarum* and its fermentation products on growth performance, immune function, intestinal pH, and cecal microorganisms of Lingnan yellow chicken. *Poultry Science*, 102(6), 102610. <https://doi.org/10.1016/j.psj.2023.102610>
- Yan, L., An, S., Lv, X., Lv, Z., Zhang, B., Choct, M., Guo, Y., Wang, Z., Yan, B., & Li, Y. (2025). Effects of replacing soybean meal with cottonseed meal on growth performance, carcass trait, intestinal development and intestinal microbiota of broiler chickens. *Poultry Science*, 104(2), 104653. <https://doi.org/10.1016/j.psj.2024.104653>
- Yang, R., Khalid, A., Khalid, F., Ye, M., Li, Y., Zhan, K., Li, Y., Liu, W., & Wang, Z. (2022). Effect of fermented corn by-products on production performance, blood biochemistry, and egg quality indices of laying hens. *Journal of Animal Science*, 100(5). <https://doi.org/10.1093/jas/skac130>
- Yu, L., Zhihui, C., Hongzhi, W., Fengjie, J., Yang, L., Jianing, L., ... & Liangmei, X. (2024). Effects of Fermented Puffed Feather Meal on Growth Performance, Serum Biochemical Indices, Meat Quality, and Intestinal Microbiota in Broilers. *Journal of Northeast Agricultural University*, 31(3).

*Thank you for publishing with us.